

Generalized Coherent States Associated with the C_λ -Extended Oscillator

C. Quesne

*Physique Nucléaire Théorique et Physique Mathématique, Université Libre de Bruxelles,
Campus de la Plaine CP229, Boulevard du Triomphe, B-1050 Brussels, Belgium*

E-mail: cquesne@ulb.ac.be

Abstract

Two new types of coherent states associated with the C_λ -extended oscillator, where C_λ is the cyclic group of order λ , are introduced. The first ones include as special cases both the Barut-Girardello and the Perelomov $\text{su}(1,1)$ coherent states for $\lambda = 2$, as well as the annihilation-operator coherent states of the C_λ -extended oscillator spectrum generating algebra for higher λ values. The second ones, which are eigenstates of the C_λ -extended oscillator annihilation operator, extend to higher λ values the paraboson coherent states, to which they reduce for $\lambda = 2$. All these states satisfy a unity resolution relation in the C_λ -extended oscillator Fock space (or in some subspace thereof). They give rise to Bargmann representations of the latter, wherein the generators of the C_λ -extended oscillator algebra are realized as differential-operator-valued matrices (or differential operators). The statistical and squeezing properties of the new coherent states are investigated over a wide range of parameters and some interesting nonclassical features are exhibited.

1 INTRODUCTION

Coherent states (CS) of the harmonic oscillator [1, 2, 3], dating back to the birth of quantum mechanics [4], are known to have properties similar to those of the classical radiation field. They are specific superpositions $|z\rangle$, parametrized by a single complex number z , of the eigenstates $|n\rangle$ of the number operator $N_b \equiv b^\dagger b$ with

$$[N_b, b^\dagger] = b^\dagger, \quad [N_b, b] = -b, \quad [b, b^\dagger] = I. \quad (1.1)$$

They may be constructed in three equivalent ways: (i) by defining them as eigenstates of the annihilation operator b ; (ii) by applying a unitary displacement operator on the vacuum state $|0\rangle$ (such that $b|0\rangle = 0$); (iii) by considering them as quantum states with a minimum uncertainty relationship.

These three construction methods become inequivalent when considering generalized CS associated with other algebras than the oscillator one, defined in (1.1). One may therefore distinguish between annihilation-operator CS (often called Barut-Girardello CS [5]), displacement-operator CS (often called Perelomov CS [6]), and minimum-uncertainty or “intelligent” CS [7], but this in no way exhausts all the possibilities of defining generalized CS.

Such states provide a fundamental framework wherein global dynamical properties of quantum systems can be interpreted [8, 9, 10]. They are commonly used for instance in connection with variational and path-integral methods or to study the classical and thermodynamic limits of quantum mechanics.

They have also found considerable applications in quantum optics, because they may exhibit some nonclassical properties, such as photon antibunching [11], or sub-Poissonian photon statistics [12], and squeezing [13]. As examples of generalized CS with nonclassical properties, we may quote the eigenstates of b^2 , which were introduced as even and odd CS or cat states [14], and are a special case of Barut-Girardello CS associated with the Lie algebra $\text{su}(1,1)$ [5]. We may also mention the eigenstates of b^λ ($\lambda > 2$) or kitten states [15], which are an example of multiphoton CS [16].

Recently there has been much interest in the study of nonlinear CS associated with a deformed oscillator (or f -oscillator). The latter is defined in terms of creation, annihilation,

and number operators, $a^\dagger = f(N_b)b^\dagger$, $a = bf(N_b)$, $N = N_b$, satisfying the commutation relations [17, 18, 19, 20]

$$[N, a^\dagger] = a^\dagger, \quad [N, a] = -a, \quad [a, a^\dagger] = G(N), \quad (1.2)$$

where f is some Hermitian operator-valued function of the number operator and $G(N) = (N+1)f^2(N+1) - Nf^2(N)$. Nonlinear CS defined as eigenstates of a [18, 20, 21, 22, 23], of a^2 [24], or of an arbitrary power a^λ ($\lambda > 2$) [25] have, for instance, been considered in connection with nonclassical properties. For a particular class of nonlinearities, the first ones were shown to be realized physically as the stationary states of the centre-of-mass motion of a trapped ion [22]. It should be noted that the algebra (1.2) may also be seen as a deformed $\text{su}(1,1)$ algebra. Hence the eigenstates of a may be considered as a deformation of the Barut-Girardello $\text{su}(1,1)$ CS [26].

The purpose of the present paper is to study some mathematical and physical properties of various CS associated with the recently introduced C_λ -extended oscillator (where $C_\lambda = \mathbb{Z}_\lambda$ denotes the cyclic group of order λ) [27]. The latter, which shares some properties with a fractional supersymmetric oscillator [28], has proved very useful in the context of supersymmetric quantum mechanics and some of its variants [27, 29, 30]. It may be considered as a deformed oscillator with a $\mathbb{Z}_\lambda$ -graded Fock space (with components labelled by $\mu = 0, 1, \dots, \lambda - 1$), generalizing the Calogero-Vasiliev [31] or modified [32] oscillator, to which it reduces for $\lambda = 2$. Its coherent states are therefore special cases of the nonlinear CS considered in [18, 20, 21, 22, 23, 24, 25, 26].

In [33], we already studied one type of CS $|z; \mu\rangle$, associated with the C_λ -extended oscillator, namely annihilation-operator CS for its spectrum generating algebra (SGA), a C_λ -extended polynomial deformation of $\text{su}(1,1)$. Since such states are eigenstates of $J_- = a^\lambda/\lambda$, corresponding to the eigenvalue z , they fall into the class of multiphoton nonlinear CS [25], and reduce to the Barut-Girardello $\text{su}(1,1)$ CS for $\lambda = 2$ and to the standard multiphoton CS [15] (equivalent to some Mittag-Leffler CS [34]) for arbitrary $\lambda > 2$ and vanishing C_λ -extended oscillator parameters. Their main mathematical feature is that they satisfy Klauder's minimal set of conditions for generalized CS [2], including the existence of a unity resolution relation, property that is not shared by generic multiphoton nonlinear CS. In addition, from a physical viewpoint, they were shown to exhibit, especially for $\mu = 0$, strong nonclassical properties, such as antibunching and quadrature squeezing.

Here we will extend the results of [33] in two different directions. Firstly, we will consider a whole family of CS, each member of the family being labelled by some index $\alpha \in \{0, 1, \dots, [\lambda/2]\}$, where $[\lambda/2]$ denotes the largest integer contained in $\lambda/2$. To each α value will be associated some CS $|z; \mu; \alpha\rangle$, $\mu = 0, 1, \dots, \lambda - \alpha - 1$, satisfying Klauder's conditions [2] in some subspace and for some appropriate values of the C_λ -extended oscillator parameters. This family will include not only the multiphoton nonlinear CS $|z, \mu\rangle$ of [33], corresponding to $\alpha = 0$, but also the Perelomov $\text{su}(1,1)$ CS [6], corresponding to $\lambda = 2$ and $\alpha = 1$.

Secondly, we will use the CS $|z; \mu; 0\rangle$, with z replaced by $\omega = z^\lambda$, to obtain the eigenstates $|z\rangle$ of the C_λ -extended oscillator annihilation operator a . For $\lambda \geq 3$, such states will generalize the paraboson CS [35], to which they will reduce for $\lambda = 2$ owing to the connection [36] between the Calogero-Vasiliev algebra [31, 32] and the paraboson one [37].

Both types of CS $|z; \mu; 0\rangle$ and $|z\rangle$ providing us with a Bargmann representation [38] of the C_λ -extended oscillator Fock space, we will use them to derive differential-operator-valued matrix realizations of the C_λ -extended oscillator algebra and of its SGA. In such a context, the μ index will play a role similar to that of the discrete label appearing in vector CS [10, 39].

This paper is organized as follows. The C_λ -extended oscillator and the corresponding algebras are briefly reviewed in Section 2. Generalized CS associated with this oscillator are introduced and proved to satisfy unity resolution relations in Sections 3 and 4, respectively. They are then used in Section 5 to construct Bargmann representations. Some physical applications are discussed in Section 6. Finally, Section 7 contains the conclusion.

2 THE C_λ -EXTENDED OSCILLATOR

The C_λ -extended oscillator algebra was introduced as a generalization of the Calogero-Vasiliev [31] or modified oscillator [32] algebra, defined by

$$[N, a^\dagger] = a^\dagger, \quad [a, a^\dagger] = I + \alpha_0 K, \quad [K, N] = \{K, a^\dagger\} = 0, \quad (2.1)$$

and their Hermitian conjugates, where α_0 is some real parameter subject to the condition $\alpha_0 > -1$, and N, K are some Hermitian operators. The latter may be realized as $K = (-1)^N$, so that the second equation in (2.1) becomes equivalent to $[a, a^\dagger] = I + \alpha_0 P_0 + \alpha_1 P_1$,

where $\alpha_0 + \alpha_1 = 0$ and $P_0 = \frac{1}{2} [I + (-1)^N]$, $P_1 = \frac{1}{2} [I - (-1)^N]$ project on the even and odd subspaces, $\mathcal{F}_0 \equiv \{|2k\rangle \mid k = 0, 1, \dots\}$, $\mathcal{F}_1 \equiv \{|2k+1\rangle \mid k = 0, 1, \dots\}$, of the Fock space $\mathcal{F} = \mathcal{F}_0 \oplus \mathcal{F}_1 = \{|n\rangle \mid n = 0, 1, \dots\}$, respectively. The oscillator Hamiltonian associated with the algebra (2.1),

$$H_0 = \frac{1}{2} \{a, a^\dagger\}, \quad (2.2)$$

provides an algebraic formulation of the two-particle Calogero problem (see [32] and references quoted therein).

When partitioning $\mathcal{F}$ into λ subspaces $\mathcal{F}_\mu \equiv \{|k\lambda + \mu\rangle \mid k = 0, 1, \dots\}$, $\mu = 0, 1, \dots, \lambda-1$, instead of two, the Calogero-Vasiliev algebra (2.1) is replaced by the C_λ -extended oscillator algebra, satisfying the relations

$$[N, a^\dagger] = a^\dagger, \quad [a, a^\dagger] = I + \sum_{\mu=0}^{\lambda-1} \alpha_\mu P_\mu, \quad [N, P_\mu] = 0, \quad a^\dagger P_\mu = P_{\mu+1} a^\dagger, \quad (2.3)$$

and their Hermitian conjugates [27, 30]. Here the α_μ 's are some real parameters subject to the conditions

$$\sum_{\mu=0}^{\lambda-1} \alpha_\mu = 0, \quad (2.4)$$

$$\sum_{\nu=0}^{\mu-1} \alpha_\nu > -\mu, \quad \mu = 1, 2, \dots, \lambda-1. \quad (2.5)$$

The operators

$$P_\mu = \frac{1}{\lambda} \sum_{\nu=0}^{\lambda-1} e^{2\pi i \nu (N-\mu)/\lambda}, \quad (2.6)$$

which are linear combinations of the operators of a cyclic group C_λ , project on the subspaces $\mathcal{F}_\mu$, and satisfy the relations $P_\mu P_\nu = \delta_{\mu,\nu} P_\mu$ and $\sum_{\mu=0}^{\lambda-1} P_\mu = I$. Throughout this paper, except otherwise stated, we use the convention $P_\mu = P_{\mu'}$ if $\mu' - \mu = 0 \bmod \lambda$ (and similarly for other operators or parameters labelled by μ, μ').

When we insert (2.6) into the second equation of (2.3), it becomes clear that the C_λ -extended oscillator algebra is a special case of deformed oscillator algebra, as defined in (1.2). Here $f(N) = [F(N)/N]^{1/2}$, where the so-called structure function $F(N)$ is given by

$$F(N) = N + \sum_{\mu=0}^{\lambda-1} \beta_\mu P_\mu, \quad \beta_\mu \equiv \sum_{\nu=0}^{\mu-1} \alpha_\nu. \quad (2.7)$$

In the Fock space representation, the operators N , $a^\dagger$, a are related to each other through the function $F(N)$,

$$a^\dagger a = F(N), \quad aa^\dagger = F(N+1). \quad (2.8)$$

The Fock space basis states

$$|n\rangle = |k\lambda + \mu\rangle = \left[\lambda^{k\lambda + \mu} k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_\nu)_{k+1} \right) \left(\prod_{\nu=\mu+1}^{\lambda-1} (\bar{\beta}_\nu)_k \right) \right]^{-1/2} (a^\dagger)^{k\lambda + \mu} |0\rangle, \quad (2.9)$$

where $n, k = 0, 1, \dots, \mu = 0, 1, \dots, \lambda - 1$, $a|0\rangle = 0$, $\bar{\beta}_\nu \equiv (\beta_\nu + \nu)/\lambda$, and $(\bar{\beta}_\nu)_k$ denotes Pochhammer's symbol, satisfy the relations

$$N|n\rangle = n|n\rangle, \quad a^\dagger|n\rangle = \sqrt{F(n+1)}|n+1\rangle, \quad a|n\rangle = \sqrt{F(n)}|n-1\rangle. \quad (2.10)$$

Due to the restriction (2.5) on the range of the parameters α_μ , $F(\mu) = \beta_\mu + \mu = \lambda\bar{\beta}_\mu > 0$ so that all the states $|n\rangle$ are well defined.

The C_λ -extended oscillator Hamiltonian is still defined by (2.2). Its eigenstates are the states $|n\rangle = |k\lambda + \mu\rangle$ and their eigenvalues are given by

$$E_{k\lambda + \mu} = k\lambda + \mu + \gamma_\mu + \frac{1}{2}, \quad \gamma_\mu \equiv \frac{1}{2}(\beta_\mu + \beta_{\mu+1}). \quad (2.11)$$

In each $\mathcal{F}_\mu$ subspace of $\mathcal{F}$, the spectrum of H_0 is harmonic, but the λ infinite sets of equally spaced energy levels, corresponding to $\mu = 0, 1, \dots, \lambda - 1$, are shifted with respect to each other by some amounts depending upon the parameters $\alpha_0, \alpha_1, \dots, \alpha_{\lambda-1}$ through their linear combinations γ_μ .

The whole spectrum of H_0 can be generated from the lowest eigenstates $|\mu\rangle$, $\mu = 0, 1, \dots, \lambda - 1$, belonging to $\mathcal{F}_\mu$, $\mu = 0, 1, \dots, \lambda - 1$, by using the operators [33]

$$J_+ = \frac{1}{\lambda} (a^\dagger)^\lambda, \quad J_- = \frac{1}{\lambda} a^\lambda, \quad J_0 = \frac{1}{\lambda} H_0 = \frac{1}{2\lambda} \{a, a^\dagger\}. \quad (2.12)$$

Equation (2.9) can indeed be rewritten as

$$|n\rangle = |k\lambda + \mu\rangle = \left[\lambda^{k(\lambda-2)} k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_\nu + 1)_k \right) \left(\prod_{\nu=\mu+1}^{\lambda-1} (\bar{\beta}_\nu)_k \right) \right]^{-1/2} J_+^k |\mu\rangle. \quad (2.13)$$

The operators (2.12) satisfy the Hermiticity properties

$$J_+^\dagger = J_-, \quad J_0^\dagger = J_0, \quad (2.14)$$

and the commutation relations

$$[J_0, J_{\pm}] = \pm J_{\pm}, \quad [J_+, J_-] = f(J_0, P_{\mu}), \quad [J_0, P_{\mu}] = [J_{\pm}, P_{\mu}] = 0, \quad (2.15)$$

where the function $f(J_0, P_{\mu})$ (not to be confused with the function $f(N)$ of (1.2)) is a $(\lambda - 1)$ th-degree polynomial in J_0 with P_{μ} -dependent coefficients, given by

$$\begin{aligned} f(J_0, P_{\mu}) &= \sum_{i=0}^{\lambda-1} s_i(P_{\mu}) J_0^i \\ &= -\frac{1}{\lambda} \left\{ \prod_{l=0}^{\lambda-2} \left(\lambda J_0 + \frac{1}{2} \sum_{\mu} \left(2l + 1 + \alpha_{\mu} + 2 \sum_{m=1}^l \alpha_{\mu+m} \right) P_{\mu} \right) \right. \\ &\quad \left. + \sum_{i=1}^{\lambda-1} \left(\lambda J_0 - \frac{1}{2} \sum_{\mu} (1 + \alpha_{\mu}) P_{\mu} \right) \right. \\ &\quad \times \left[\prod_{j=1}^{i-1} \left(\lambda J_0 + \frac{1}{2} \sum_{\mu} \left(-2j - 1 + \alpha_{\mu} + 2 \sum_{k=1}^{\lambda-j-1} \alpha_{\mu+k} \right) P_{\mu} \right) \right] \\ &\quad \times \left[\prod_{l=0}^{\lambda-i-2} \left(\lambda J_0 + \frac{1}{2} \sum_{\mu} \left(2l + 1 + \alpha_{\mu} + 2 \sum_{m=1}^l \alpha_{\mu+m} \right) P_{\mu} \right) \right] \left. \right\}. \quad (2.16) \end{aligned}$$

The SGA of the C_{λ} -extended oscillator is therefore a C_{λ} -extended polynomial deformation of the Lie algebra $\mathfrak{su}(1,1)$: in each $\mathcal{F}_{\mu}$ subspace or for vanishing algebra parameters, it reduces to a standard polynomial deformation of $\mathfrak{su}(1,1)$ [40]. Nonlinearities are only present for $\lambda \geq 3$, since for $\lambda = 2$ the algebra reduces to the well-known $\mathfrak{su}(1,1)$ SGA of the Calogero-Vasiliev oscillator [32].

The algebra (2.14)–(2.16) has a Casimir operator, which can be written as

$$C = J_- J_+ + h(J_0, P_{\mu}) = J_+ J_- + h(J_0, P_{\mu}) - f(J_0, P_{\mu}), \quad (2.17)$$

where $h(J_0, P_{\mu})$ is a λ th-degree polynomial in J_0 with P_{μ} -dependent coefficients, $h(J_0, P_{\mu}) = \sum_{i=0}^{\lambda} t_i(P_{\mu}) J_0^i$, whose explicit form is given in [33].

Each $\mathcal{F}_{\mu}$ subspace of $\mathcal{F}$ carries a unitary irreducible representation (unirrep) of the SGA, characterized by the eigenvalue c_{μ} of C and by the lowest eigenvalue of J_0 , namely $(\mu + \gamma_{\mu} + \frac{1}{2})/\lambda$.

3 GENERALIZED COHERENT STATES ASSOCIATED WITH THE C_{λ} -EXTENDED OSCILLATOR

3.1 Family of Coherent States $|z; \mu; \alpha\rangle$

Let $|z; \mu; \alpha\rangle$ be a normalizable state of $\mathcal{F}_\mu$, satisfying the equation

$$\left[a^{\lambda-\alpha} - z(a^\dagger)^\alpha\right] |z; \mu; \alpha\rangle = 0 \quad (3.1)$$

for some $\alpha \in \{0, 1, \dots, [\lambda/2]\}$ and some $z \in \mathbb{C}$. The set of states $|z; \mu; \alpha\rangle$ is surely non-empty, since for $\alpha = 0$, Eq. (3.1) reduces to

$$a^\lambda |z; \mu; 0\rangle = z |z; \mu; 0\rangle, \quad (3.2)$$

showing that $|z; \mu; 0\rangle$ is an annihilation-operator CS $|z/\lambda; \mu\rangle$ for the C_λ -extended oscillator SGA, corresponding to the eigenvalue z/λ of J_- ,

$$|z; \mu; 0\rangle = |z/\lambda; \mu\rangle. \quad (3.3)$$

The latter is known to be normalizable for any $z \in \mathbb{C}$ and any $\mu \in \{0, 1, \dots, \lambda - 1\}$ [33].

Furthermore, for $\lambda = 2$, $\mu = 0$, and $\alpha = 1$, Eq. (3.1) becomes

$$(a - za^\dagger) |z; 0; 1\rangle = 0. \quad (3.4)$$

By using Baker-Campbell-Hausdorff formula, this equation can be rewritten as

$$e^{zJ_+} a e^{-zJ_+} |z; 0; 1\rangle = 0, \quad (3.5)$$

where J_+ is given by (2.12) where $\lambda = 2$. It is therefore clear that

$$|z; 0; 1\rangle = (1 - |z|^2)^{\bar{\beta}_1/2} e^{zJ_+} |0\rangle \quad (3.6)$$

is a displacement-operator (or Perelomov [6]) CS for $\mathfrak{su}(1,1)$, corresponding to the unirrep characterized by $\frac{1}{2}(\gamma_0 + \frac{1}{2}) = \frac{1}{2}\bar{\beta}_1$. Such a state is defined on the unit disc $|z| < 1$.

We may therefore say that the states (3.1) in some way interpolate between Barut-Girardello and Perelomov CS. We shall now proceed to construct them and to determine under which conditions they are normalizable.

Let us expand $|z; \mu; \alpha\rangle$ in the number-state basis $\{|k\lambda + \mu\rangle \mid k = 0, 1, \dots\}$ of $\mathcal{F}_\mu$,

$$|z; \mu; \alpha\rangle = \sum_{k=0}^{\infty} c_k(z, z^*; \mu; \alpha) z^k |k\lambda + \mu\rangle, \quad (3.7)$$

where $c_k(z, z^*; \mu; \alpha)$ are some yet unknown coefficients. By iterating (2.10) and taking (2.7) into account, we easily obtain

$$\begin{aligned}
a^{\lambda-\alpha} |k\lambda + \mu\rangle &= \left[\lambda^{\lambda-\alpha} k \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_{\nu} + k) \right) \left(\prod_{\nu=\mu+\alpha+1}^{\lambda-1} (\bar{\beta}_{\nu} + k - 1) \right) \right]^{1/2} \\
&\quad \times |(k-1)\lambda + \mu + \alpha\rangle \quad \text{if } \mu = 0, 1, \dots, \lambda - \alpha - 1, \\
&= \left[\lambda^{\lambda-\alpha} \left(\prod_{\nu=\mu-\lambda+\alpha+1}^{\mu} (\bar{\beta}_{\nu} + k) \right) \right]^{1/2} |k\lambda + \mu - \lambda + \alpha\rangle \\
&\quad \text{if } \mu = \lambda - \alpha, \lambda - \alpha + 1, \dots, \lambda - 1,
\end{aligned} \tag{3.8}$$

and

$$\begin{aligned}
(a^\dagger)^\alpha |k\lambda + \mu\rangle &= \left[\lambda^\alpha \left(\prod_{\nu=\mu+1}^{\mu+\alpha} (\bar{\beta}_{\nu} + k) \right) \right]^{1/2} |k\lambda + \mu + \alpha\rangle \quad \text{if } \mu = 0, 1, \dots, \lambda - \alpha - 1, \\
&= \left[\lambda^\alpha (k+1) \left(\prod_{\nu=1}^{\mu-\lambda+\alpha} (\bar{\beta}_{\nu} + k + 1) \right) \left(\prod_{\nu=\mu+1}^{\lambda-1} (\bar{\beta}_{\nu} + k) \right) \right]^{1/2} \\
&\quad \times |(k+1)\lambda + \mu - \lambda + \alpha\rangle \quad \text{if } \mu = \lambda - \alpha, \lambda - \alpha + 1, \dots, \lambda - 1.
\end{aligned} \tag{3.9}$$

Inserting (3.7), (3.8), and (3.9) into (3.1) leads to a recursion relation for $c_k(z, z^*; \mu; \alpha)$. For $\mu \in \{0, 1, \dots, \lambda - \alpha - 1\}$, the latter has a solution given by

$$c_k(z, z^*; \mu; \alpha) = \left(\frac{\lambda^{k(2\alpha-\lambda)} \left(\prod_{\nu=\mu+1}^{\mu+\alpha} (\bar{\beta}_{\nu})_k \right)}{k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_{\nu} + 1)_k \right) \left(\prod_{\nu=\mu+\alpha+1}^{\lambda-1} (\bar{\beta}_{\nu})_k \right)} \right)^{1/2} c_0(z, z^*; \mu; \alpha), \tag{3.10}$$

while for $\mu \in \{\lambda - \alpha, \lambda - \alpha + 1, \dots, \lambda - 1\}$, it has no solution apart from the trivial one, $c_k(z, z^*; \mu; \alpha) = 0$, $k = 0, 1, \dots$.

By introducing (3.10) into (3.7), we obtain

$$\begin{aligned}
|z; \mu; \alpha\rangle &= \left[N_\mu^{(\alpha)}(|z|) \right]^{-1/2} \sum_{k=0}^{\infty} \left(\frac{\prod_{\nu=\mu+1}^{\mu+\alpha} (\bar{\beta}_{\nu})_k}{k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_{\nu} + 1)_k \right) \left(\prod_{\nu=\mu+\alpha+1}^{\lambda-1} (\bar{\beta}_{\nu})_k \right)} \right)^{1/2} \\
&\quad \times \left(\lambda^{-(\lambda-2\alpha)/2} z \right)^k |k\lambda + \mu\rangle,
\end{aligned} \tag{3.11}$$

where the normalization coefficient $N_\mu^{(\alpha)}(|z|)$ can be expressed in terms of a generalized hypergeometric function,

$$\begin{aligned}
N_\mu^{(\alpha)}(|z|) &\equiv [c_0(z, z^*; \mu; \alpha)]^{-2} \\
&= {}_\alpha F_{\lambda-\alpha-1} \left(\bar{\beta}_{\mu+1}, \dots, \bar{\beta}_{\mu+\alpha}; \bar{\beta}_1 + 1, \dots, \bar{\beta}_\mu + 1, \bar{\beta}_{\mu+\alpha+1}, \dots, \bar{\beta}_{\lambda-1}; y \right), \\
y &\equiv \frac{|z|^2}{\lambda^{\lambda-2\alpha}}.
\end{aligned} \tag{3.12}$$

Since the latter converges for all finite y if $\alpha \leq \lambda - \alpha - 1$, and for $|y| < 1$ if $\alpha = \lambda - \alpha$, we conclude that Eq. (3.1) has normalizable solutions defined on the complex plane for any $\alpha \in \{0, 1, \dots, [(\lambda - 1)/2]\}$ and any $\mu \in \{0, 1, \dots, \lambda - \alpha - 1\}$, and normalizable solutions defined on the unit disc for λ even, $\alpha = \lambda/2$, and any $\mu \in \{0, 1, \dots, (\lambda - 2)/2\}$.

In the former subset, there are the CS (3.3) of the C_λ -extended oscillator SGA. As new examples, we find, for instance, for $\lambda = 3$ the states

$$|z; 0; 1\rangle = [{}_1F_1(\bar{\beta}_1; \bar{\beta}_2; y)]^{-1/2} \sum_{k=0}^{\infty} \left(\frac{(\bar{\beta}_1)_k}{k! (\bar{\beta}_2)_k} \right)^{1/2} \left(\frac{z}{\sqrt{3}} \right)^k |3k\rangle, \quad (3.13)$$

$$|z; 1; 1\rangle = [{}_1F_1(\bar{\beta}_2; \bar{\beta}_1 + 1; y)]^{-1/2} \sum_{k=0}^{\infty} \left(\frac{(\bar{\beta}_2)_k}{k! (\bar{\beta}_1 + 1)_k} \right)^{1/2} \left(\frac{z}{\sqrt{3}} \right)^k |3k + 1\rangle, \quad (3.14)$$

with $y \equiv |z|^2/3$, which satisfy the equation $(a^2 - za^\dagger)|z; \mu; 1\rangle = 0$, $\mu = 0, 1$.

The Perelomov $\text{su}(1,1)$ CS (3.6) belong to the latter subset, wherein we also encounter, for instance, for $\lambda = 4$ the states

$$|z; 0; 2\rangle = [{}_2F_1(\bar{\beta}_1, \bar{\beta}_2; \bar{\beta}_3; y)]^{-1/2} \sum_{k=0}^{\infty} \left(\frac{(\bar{\beta}_1)_k (\bar{\beta}_2)_k}{k! (\bar{\beta}_3)_k} \right)^{1/2} z^k |4k\rangle, \quad (3.15)$$

$$|z; 1; 2\rangle = [{}_2F_1(\bar{\beta}_2, \bar{\beta}_3; \bar{\beta}_1 + 1; y)]^{-1/2} \sum_{k=0}^{\infty} \left(\frac{(\bar{\beta}_2)_k (\bar{\beta}_3)_k}{k! (\bar{\beta}_1 + 1)_k} \right)^{1/2} z^k |4k + 1\rangle, \quad (3.16)$$

with $y \equiv |z|^2$, which are solutions of $[a^2 - z(a^\dagger)^2]|z; \mu; 2\rangle = 0$, $\mu = 0, 1$.

By using (2.13), Eq. (3.11) can be written in an alternative form

$$|z; \mu; \alpha\rangle = [N_\mu^{(\alpha)}(|z|)]^{-1/2} {}_0F_{\lambda-\alpha-1}(\bar{\beta}_1 + 1, \dots, \bar{\beta}_\mu + 1, \bar{\beta}_{\mu+\alpha+1}, \dots, \bar{\beta}_{\lambda-1}; zJ_+/\lambda^{\lambda-\alpha-1}) |\mu\rangle \quad (3.17)$$

in terms of the SGA generator J_+ .

The overlap of two states of type (3.11) can be easily calculated and is given by

$$\begin{aligned} \langle z'; \mu'; \alpha' | z; \mu; \alpha \rangle &= \delta_{\mu', \mu} [N_\mu^{(\alpha)}(|z|) N_\mu^{(\alpha')}(|z|)]^{-1/2} \\ &\times {}_{\alpha_m}F_{\lambda-\alpha_M-1}(\bar{\beta}_{\mu+1}, \dots, \bar{\beta}_{\mu+\alpha_m}; \bar{\beta}_1 + 1, \dots, \bar{\beta}_\mu + 1, \bar{\beta}_{\mu+\alpha_M+1}, \dots, \bar{\beta}_{\lambda-1}; \\ &\quad z'^* z / \lambda^{\lambda-\alpha-\alpha'}), \end{aligned} \quad (3.18)$$

where $\alpha_m \equiv \min(\alpha, \alpha')$ and $\alpha_M \equiv \max(\alpha, \alpha')$. From the continuity of the overlapping factor $\langle z'; \mu; \alpha | z; \mu; \alpha \rangle$, it follows that the normalizable states $|z; \mu; \alpha\rangle$ are continuous in the label z , i.e.,

$$|z - z'| \rightarrow 0 \quad \Rightarrow \quad \left| |z; \mu; \alpha\rangle - |z'; \mu; \alpha\rangle \right|^2 \rightarrow 0. \quad (3.19)$$

Hence, they satisfy two of the three conditions considered by Klauder [2] as minimal for the existence of generalized CS. We leave the discussion of the third one, i.e., the existence of a unity resolution relation, for the next Section.

3.2 Eigenstates $|z\rangle$ of the Annihilation Operator a

Let us now look for the eigenstates $|z\rangle$ of the C_λ -extended oscillator annihilation operator a ,

$$a|z\rangle = z|z\rangle, \quad (3.20)$$

corresponding to some eigenvalues $z \in \mathbb{C}$.

We shall try to construct them as linear combinations of the CS $|\omega; \mu; 0\rangle$, $\mu = 0, 1, \dots, \lambda - 1$, considered in the previous Subsection, where we replace z by $\omega \equiv z^\lambda$,

$$|z\rangle = \sum_{\mu=0}^{\lambda-1} d_\mu(z, z^*) \left(\frac{z}{\sqrt{\lambda}} \right)^\mu |\omega; \mu; 0\rangle. \quad (3.21)$$

Here $d_\mu(z, z^*)$ are some coefficients to be determined.

From (2.7), (2.10), (3.7), and (3.10), we easily obtain

$$a|\omega; 0; 0\rangle = \left[\lambda^{\lambda-1} \left(\prod_{\nu=1}^{\lambda-1} \bar{\beta}_\nu \right) \right]^{-1/2} \frac{c_0(\omega, \omega^*; 0; 0)}{c_0(\omega, \omega^*; \lambda-1; 0)} \omega |\omega; \lambda-1; 0\rangle, \quad (3.22)$$

$$a|\omega; \mu; 0\rangle = \sqrt{\lambda \bar{\beta}_\mu} \frac{c_0(\omega, \omega^*; \mu; 0)}{c_0(\omega, \omega^*; \mu-1; 0)} |\omega; \mu-1; 0\rangle, \quad \mu = 1, 2, \dots, \lambda-1. \quad (3.23)$$

Inserting (3.21), (3.22), and (3.23) into (3.20) leads to

$$d_\mu(z, z^*) = \left(\prod_{\nu=1}^{\mu} \bar{\beta}_\nu \right)^{-1/2} \frac{c_0(\omega, \omega^*; 0; 0)}{c_0(\omega, \omega^*; \mu; 0)} d_0(z, z^*), \quad \mu = 1, 2, \dots, \lambda-1. \quad (3.24)$$

By taking (3.12) into account, the solutions of (3.20) can therefore be written as

$$|z\rangle = [\mathcal{N}(|z|)]^{-1/2} \sum_{\mu=0}^{\lambda-1} \left(\frac{N_\mu^{(0)}(|\omega|)}{\prod_{\nu=1}^{\mu} \bar{\beta}_\nu} \right)^{1/2} \left(\frac{z}{\sqrt{\lambda}} \right)^\mu |\omega; \mu; 0\rangle, \quad (3.25)$$

where the normalization factor $\mathcal{N}(|z|)$ is given by a linear combination of generalized hypergeometric functions,

$$\begin{aligned} \mathcal{N}(|z|) &\equiv [c_0(z, z^*; 0; 0) d_0(z, z^*)]^{-2} \\ &= \sum_{\mu=0}^{\lambda-1} N_\mu^{(0)}(|z|^\lambda) \frac{(|z|^2/\lambda)^\mu}{\prod_{\nu=1}^{\mu} \bar{\beta}_\nu} \\ &= \sum_{\mu=0}^{\lambda-1} {}_0F_{\lambda-1}(\bar{\beta}_1 + 1, \dots, \bar{\beta}_\mu + 1, \bar{\beta}_{\mu+1}, \dots, \bar{\beta}_{\lambda-1}; t^\lambda) \frac{t^\mu}{\prod_{\nu=1}^{\mu} \bar{\beta}_\nu}, \end{aligned} \quad (3.26)$$

with $t \equiv |z|^2/\lambda$. As the CS $|\omega; \mu; 0\rangle$, the annihilation operator eigenstates $|z\rangle$ are defined on the whole complex plane.

By using (3.11), they can be rewritten in the number-state basis $\{|k\lambda + \mu\rangle \mid k = 0, 1, \dots, \mu = 0, 1, \dots, \lambda - 1\}$ of $\mathcal{F}$ as

$$|z\rangle = [\mathcal{N}(|z|)]^{-1/2} \sum_{\mu=0}^{\lambda-1} \sum_{k=0}^{\infty} \frac{(z/\sqrt{\lambda})^{k\lambda+\mu}}{\left[k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_{\nu})_{k+1}\right) \left(\prod_{\nu=\mu+1}^{\lambda-1} (\bar{\beta}_{\nu})_k\right)\right]^{1/2}} |k\lambda + \mu\rangle. \quad (3.27)$$

From (3.18) and (3.25), the overlap of two eigenstates of a is obtained as

$$\begin{aligned} \langle z'|z\rangle &= [\mathcal{N}(|z|)\mathcal{N}(|z'|)]^{-1/2} \sum_{\mu=0}^{\lambda-1} {}_0F_{\lambda-1}(\bar{\beta}_1 + 1, \dots, \bar{\beta}_{\mu} + 1, \bar{\beta}_{\mu+1}, \dots, \bar{\beta}_{\lambda-1}; (z'^* z/\lambda)^{\lambda}) \\ &\quad \times \frac{(z'^* z/\lambda)^{\mu}}{\prod_{\nu=1}^{\mu} \bar{\beta}_{\nu}}, \end{aligned} \quad (3.28)$$

which is a continuous function. Hence the normalizable states $|z\rangle$ are continuous in z , i.e., satisfy a property similar to (3.19).

Before proceeding to the determination of a unity resolution relation in the next Section, it is worth considering the example $\lambda = 2$, corresponding to the Calogero-Vasiliev algebra [31, 32] or, equivalently, to the paraboson one [37]. In such a case, the ${}_0F_1$ function appearing in (3.26) and (3.28) can be expressed in terms of a modified Bessel function I_{ν} . Hence we get

$$\begin{aligned} |z\rangle &= [\mathcal{N}(|z|)]^{-1/2} \left\{ [N_0^{(0)}(|z|^2)]^{1/2} |z^2; 0; 0\rangle + [N_1^{(0)}(|z|^2)]^{1/2} \frac{z}{\sqrt{2\bar{\beta}_1}} |z^2; 1; 0\rangle \right\} \\ &= \left(\frac{\Gamma(\bar{\beta}_1)}{\mathcal{N}(|z|)} \right)^{1/2} \sum_{n=0}^{\infty} \frac{(z/\sqrt{2})^n}{\{[n/2]! \Gamma(\bar{\beta}_1 + [(n+1)/2])\}^{1/2}} |n\rangle, \end{aligned} \quad (3.29)$$

where

$$N_{\mu}^{(0)}(|z|^2) = \Gamma(\bar{\beta}_1 + \mu) \left(|z|^2/2\right)^{1-\bar{\beta}_1-\mu} I_{\bar{\beta}_1-1+\mu}(|z|^2), \quad \mu = 0, 1, \quad (3.30)$$

and

$$\mathcal{N}(|z|) = \Gamma(\bar{\beta}_1) t^{1-\bar{\beta}_1} [I_{\bar{\beta}_1-1}(2t) + I_{\bar{\beta}_1}(2t)], \quad t \equiv |z|^2/2. \quad (3.31)$$

These results coincide with those previously obtained in [35] for paraboson CS.

4 UNITY RESOLUTION RELATIONS

4.1 Unity Resolution Relations for the Coherent States $|z; \mu; \alpha\rangle$

4.1.1 Unity Resolution Relations in $\mathcal{F}_\mu$

Let us first restrict ourselves to a given subspace $\mathcal{F}_\mu$ of the Fock space $\mathcal{F}$. In such a subspace, the number states $|k\lambda + \mu\rangle$, $k = 0, 1, \dots$, form a complete set and their resolution of unity is given by a sum of orthogonal projection operators, $\sum_{k=0}^{\infty} |k\lambda + \mu\rangle\langle k\lambda + \mu| = I_\mu$, where I_μ denotes the unit operator in $\mathcal{F}_\mu$.

For any $\alpha \leq \lambda - \mu - 1$, the CS $|z; \mu; \alpha\rangle$ give a resolution of unity in $\mathcal{F}_\mu$ with a positive measure $d\rho_\mu^{(\alpha)}(z, z^*)$ if

$$\int d\rho_\mu^{(\alpha)}(z, z^*) |z; \mu; \alpha\rangle\langle z; \mu; \alpha| = I_\mu, \quad (4.1)$$

where the integration is carried out over the complex plane or the unit disc according to whether $\alpha \leq [(\lambda - 1)/2]$ or $\alpha = \lambda/2$ and λ even. Making a polar decomposition $z = |z| \exp(i\phi)$ and the ansatz

$$\begin{aligned} d\rho_\mu^{(\alpha)}(z, z^*) &= {}_\alpha F_{\lambda-\alpha-1}(\bar{\beta}_{\mu+1}, \dots, \bar{\beta}_{\mu+\alpha}; \bar{\beta}_1 + 1, \dots, \bar{\beta}_\mu + 1, \bar{\beta}_{\mu+\alpha+1}, \dots, \bar{\beta}_{\lambda-1}; y) \\ &\times h_\mu^{(\alpha)}(y) d^2z, \quad y \equiv |z|^2/\lambda^{\lambda-2\alpha}, \quad d^2z \equiv |z| d|z| d\phi, \end{aligned} \quad (4.2)$$

where $h_\mu^{(\alpha)}(y)$ is a yet unknown weight function, we find that Eq. (4.1) is equivalent to

$$\begin{aligned} \pi \lambda^{\lambda-2\alpha} \sum_{k=0}^{\infty} &\left(\frac{\prod_{\nu=\mu+1}^{\mu+\alpha} (\bar{\beta}_\nu)_k}{k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_\nu + 1)_k \right) \left(\prod_{\nu=\mu+\alpha+1}^{\lambda-1} (\bar{\beta}_\nu)_k \right)} \int_0^{y_{max}} dy y^k h_\mu^{(\alpha)}(y) \right) \\ &\times |k\lambda + \mu\rangle\langle k\lambda + \mu| = I_\mu, \end{aligned} \quad (4.3)$$

where $y_{max} = \infty$ for $\alpha \leq [(\lambda - 1)/2]$ and $y_{max} = 1$ for $\alpha = \lambda/2$ and λ even. Consequently, the requirement that for a given α , $|z; \mu; \alpha\rangle$ form a complete (in fact, an overcomplete) set in $\mathcal{F}_\mu$ is equivalent to the resolution of a Stieltjes or a Hausdorff power-moment problem [41], namely determine a positive weight function $h_\mu^{(\alpha)}(y)$ such that

$$\begin{aligned} \int_0^{y_{max}} dy y^k h_\mu^{(\alpha)}(y) &= A_\mu^{(\alpha)} \frac{\Gamma(k+1) \left(\prod_{\nu=1}^{\mu} \Gamma(\bar{\beta}_\nu + k + 1) \right) \left(\prod_{\nu=\mu+\alpha+1}^{\lambda-1} \Gamma(\bar{\beta}_\nu + k) \right)}{\prod_{\nu=\mu+1}^{\mu+\alpha} \Gamma(\bar{\beta}_\nu + k)} \\ &\equiv B_\mu^{(\alpha)}(k), \quad k = 0, 1, 2, \dots, \end{aligned} \quad (4.4)$$

where

$$A_\mu^{(\alpha)} \equiv \frac{\prod_{\nu=\mu+1}^{\mu+\alpha} \Gamma(\bar{\beta}_\nu)}{\pi \lambda^{\lambda-2\alpha} \left(\prod_{\nu=1}^{\mu} \Gamma(\bar{\beta}_\nu + 1) \right) \left(\prod_{\nu=\mu+\alpha+1}^{\lambda-1} \Gamma(\bar{\beta}_\nu) \right)} \quad (4.5)$$

is a positive constant.

The general theory of the power-moment problem [41] provides us with conditions for the solvability of (4.4). In the case where $y_{max} = \infty$, for instance, what is required is the positivity of the two series $\{H_0^{(k)}\}, \{H_1^{(k)}\}$ ($k = 1, 2, 3, \dots$) of so-called Hankel-Hadamard matrices, defined by

$$H_0^{(k)}(i, j) = B_\mu^{(\alpha)}(i + j - 2), \quad H_1^{(k)}(i, j) = B_\mu^{(\alpha)}(i + j - 1), \quad i, j = 1, 2, \dots, k. \quad (4.6)$$

The proof of positivity being extremely difficult, we will instead directly construct a solution of (4.4) in both cases $y_{max} = \infty$ and $y_{max} = 1$. For such a purpose, we will interpret the latter as an inverse Mellin transform problem [42], as done in related contexts before [23, 26, 33, 34, 43, 44], by setting for complex s , $k \rightarrow s - 1$ and rewriting (4.4) as

$$\int_0^\infty dy y^{s-1} g(y) = \frac{\prod_{\nu=1}^{r+\alpha} \Gamma(s + b_\nu)}{\prod_{\nu=1}^\alpha \Gamma(s + a_\nu)} \equiv g^*(s), \quad (4.7)$$

where for simplicity's sake we skipped the (α) and μ indices. The function $g(y) \equiv (A_\mu^{(\alpha)})^{-1} h_\mu^{(\alpha)}(y)$, which is the inverse Mellin transform of $g^*(s)$, is assumed to vanish for $y > 1$ whenever $y_{max} = 1$, and r, a_ν, b_ν are defined by

$$r = \lambda - 2\alpha \geq 0, \quad (4.8)$$

$$a_\nu = \bar{\beta}_{\mu+\nu} - 1, \quad \nu = 1, 2, \dots, \alpha, \quad (4.9)$$

$$b_\nu = \begin{cases} 0, & \nu = 1, \\ \bar{\beta}_{\nu-1}, & \nu = 2, 3, \dots, \mu + 1, \\ \bar{\beta}_{\nu+\alpha-1} - 1, & \nu = \mu + 2, \mu + 3, \dots, r + \alpha. \end{cases} \quad (4.10)$$

It is worth noting that from the restriction (2.5) and the definition of $\bar{\beta}_\nu$ given below (2.9), it results that the parameters a_ν and b_ν are greater than -1 . In addition, $r > 0$ corresponds to $y_{max} = \infty$, whereas $r = 0$ is associated with $y_{max} = 1$.

To ensure the positivity of $g(y)$ on $(0, y_{max})$, and hence that of $h_\mu^{(\alpha)}(y)$, we shall proceed step by step by using the Mellin convolution property of inverse Mellin transforms [42], which states that if $g^*(s) = g_1^*(s)g_2^*(s)$ and $g_1(y), g_2(y)$ exist, then the inverse Mellin transform of $g^*(s)$ is

$$g(y) = \int_0^\infty g_1\left(\frac{y}{t}\right) g_2(t) \frac{dt}{t} = \int_0^\infty g_1(t) g_2\left(\frac{y}{t}\right) \frac{dt}{t}. \quad (4.11)$$

In the present context, if we are sure that $g_1(y)$ and $g_2(y)$ are positive on $(0, y_{max})$, then the same will be true for $g(y)$.

For $r > 0$ and $\alpha = 0$, it is known [23, 26, 33] that Eq. (4.7) has a positive solution in terms of a Meijer G -function [45],

$$g(y) = G_{0r}^{r0}(y|b_1, b_2, \dots, b_r), \quad (4.12)$$

for all allowed values of b_ν . In Appendix A, we use this result as the starting point of an induction process over α , based upon (4.11), to show that for any $r > 0$ and any $0 \leq \alpha \leq \lambda - \mu - 1$, Eq. (4.7) has a positive solution, given by

$$g(y) = G_{\alpha, r+\alpha}^{r+\alpha, 0} \left(y \left| \begin{array}{c} a_1, a_2, \dots, a_\alpha \\ b_1, b_2, \dots, b_{r+\alpha} \end{array} \right. \right), \quad (4.13)$$

if the set $\{b_1, b_2, \dots, b_{r+\alpha}\}$ contains some subset of α elements $\{b_{i_1}, b_{i_2}, \dots, b_{i_\alpha}\}$ such that

$$a_1 > b_{i_1}, \quad a_2 > b_{i_2}, \quad \dots, \quad a_\alpha > b_{i_\alpha}. \quad (4.14)$$

We conclude that for any $\alpha \leq \min(\lambda - \mu - 1, [(\lambda - 1)/2])$, there is a positive weight function $h_\mu^{(\alpha)}(y)$, solution of (4.4) and given by

$$h_\mu^{(\alpha)}(y) = A_\mu^{(\alpha)} G_{\alpha, \lambda-\alpha}^{\lambda-\alpha, 0} \left(y \left| \begin{array}{c} \bar{\beta}_{\mu+1} - 1, \dots, \bar{\beta}_{\mu+\alpha} - 1 \\ 0, \bar{\beta}_1, \dots, \bar{\beta}_\mu, \bar{\beta}_{\mu+\alpha+1} - 1, \dots, \bar{\beta}_{\lambda-1} - 1 \end{array} \right. \right), \quad (4.15)$$

if condition (4.14) is satisfied by the sets $\{a_1, a_2, \dots, a_\alpha\}$ and $\{b_1, b_2, \dots, b_{r+\alpha}\}$ defined in (4.9) and (4.10), respectively.

As an example, let us consider the case of $\mathcal{F}_0$ and of the CS $|z; 0; 1\rangle$ corresponding to $\lambda = 3$, $\alpha = 1$, defined in (3.13). From the above results, it follows that if $\bar{\beta}_1 > 1$ or $\bar{\beta}_1 > \bar{\beta}_2$ (i.e., $\alpha_0 > 2$ or $\alpha_1 < -1$), they satisfy the unity resolution relation (4.1) with

$$h_0^{(1)}(y) = \frac{\Gamma(\bar{\beta}_1)}{3\pi\Gamma(\bar{\beta}_2)} G_{12}^{20} \left(y \left| \begin{array}{c} \bar{\beta}_1 - 1 \\ 0, \bar{\beta}_2 - 1 \end{array} \right. \right) = \frac{\Gamma(\bar{\beta}_1)}{3\pi\Gamma(\bar{\beta}_2)} e^{-y} U(\bar{\beta}_1 - \bar{\beta}_2, 2 - \bar{\beta}_2, y), \quad (4.16)$$

where $U(a, b, z) = \Psi(a, b; z)$ is Kummer's confluent hypergeometric function [45]. As illustrated in Fig. 1, $h_0^{(1)}(y) \rightarrow 0$ for $y \rightarrow \infty$ as it should be, and for $y \rightarrow 0$, $h_0^{(1)}(y) \rightarrow +\infty$ or $(\bar{\beta}_1 - 1)[3\pi(\bar{\beta}_2 - 1)]^{-1}$ according to whether $\bar{\beta}_2 \leq 1$ or $\bar{\beta}_2 > 1$ (i.e., $\alpha_0 + \alpha_1 \leq 1$ or $\alpha_0 + \alpha_1 > 1$).

It now remains to consider the case where $r = 0$, or $\alpha = \lambda/2$, λ even (and $\mu \leq \alpha - 1$). For $\lambda = 2$, $\alpha = 1$ (and $\mu = 0$), corresponding to the Perelomov $\text{su}(1,1)$ CS $|z; 0; 1\rangle$, defined

in (3.6), it is well known that there exists a positive weight function $h_0^{(1)}(y)$ on $(0, 1)$, solution of (4.4) and given by [9]

$$h_0^{(1)}(y) = \frac{\bar{\beta}_1 - 1}{\pi} (1 - y)^{\bar{\beta}_1 - 2} = \frac{\bar{\beta}_1 - 1}{\pi} {}_1F_0(2 - \bar{\beta}_1; y), \quad (4.17)$$

provided the condition $\bar{\beta}_1 > 1$ (i.e., $\alpha_0 > 1$) is satisfied. The problem to be solved is therefore finding a solution of (4.7), positive on $(0, 1)$ and vanishing on $(1, \infty)$, for $r = 0$ and $\alpha \geq 2$.

By using the convolution property (4.11) again, it is shown in Appendix B that for $r = 0$ and $\alpha = 2$, Eq. (4.7) has indeed a positive solution (B.6) on $(0, 1)$ (vanishing on $(1, \infty)$) if either $a_1 > b_1$, $a_2 > b_2$ or $a_1 > b_2$, $a_2 > b_1$. It leads to the following result for the weight function $h_0^{(2)}(y)$ of the CS $|z; 0; 2\rangle$, defined in (3.15),

$$h_0^{(2)}(y) = A_0^{(2)} \frac{(1 - y)^{\bar{\beta}_1 + \bar{\beta}_2 - \bar{\beta}_3 - 2}}{\Gamma(\bar{\beta}_1 + \bar{\beta}_2 - \bar{\beta}_3 - 1)} {}_2F_1\left(\bar{\beta}_1 - \bar{\beta}_3, \bar{\beta}_2 - \bar{\beta}_3; \bar{\beta}_1 + \bar{\beta}_2 - \bar{\beta}_3 - 1; 1 - y\right), \quad (4.18)$$

if either $\bar{\beta}_1 > \bar{\beta}_3$, $\bar{\beta}_2 > 1$ or $\bar{\beta}_1 > 1$, $\bar{\beta}_2 > \bar{\beta}_3$ (i.e., $\alpha_0 + \alpha_1 > 2$, $\alpha_1 + \alpha_2 < -2$ or $\alpha_0 > 3$, $\alpha_2 < -1$). The weight function $h_1^{(2)}(y)$ of the CS $|z; 1; 2\rangle$, defined in (3.16), can be obtained from the latter by performing the substitutions $\bar{\beta}_1 \rightarrow \bar{\beta}_2$, $\bar{\beta}_2 \rightarrow \bar{\beta}_3$, $\bar{\beta}_3 \rightarrow \bar{\beta}_1 + 1$. As illustrated in Fig. 2, for $y \rightarrow 0$, $h_0^{(2)}(y) \rightarrow +\infty$ or $(\bar{\beta}_1 - 1)(\bar{\beta}_2 - 1)[\pi(\bar{\beta}_3 - 1)]^{-1}$ according to whether $\bar{\beta}_3 \leq 1$ or $\bar{\beta}_3 > 1$ (i.e., $\alpha_0 + \alpha_1 + \alpha_2 \leq 1$ or $\alpha_0 + \alpha_1 + \alpha_2 > 1$), and for $y \rightarrow 1$, $h_0^{(2)}(y) \rightarrow +\infty$, $\Gamma(\bar{\beta}_1)\Gamma(\bar{\beta}_2)[\pi\Gamma(\bar{\beta}_3)]^{-1}$, or 0 according to whether $\bar{\beta}_1 + \bar{\beta}_2 - \bar{\beta}_3 < 2$, $\bar{\beta}_1 + \bar{\beta}_2 - \bar{\beta}_3 = 2$, or $\bar{\beta}_1 + \bar{\beta}_2 - \bar{\beta}_3 > 2$ (i.e., $\alpha_0 - \alpha_2 < 8$, $\alpha_0 - \alpha_2 = 8$, or $\alpha_0 - \alpha_2 > 8$).

For $r = 0$ and $\alpha > 2$, Eq. (4.18) can be generalized in the form of a formal multiple series of hypergeometric functions (see Appendix B). For $\mu = 0$, the result can be written as

$$\begin{aligned} h_0^{(\alpha)}(y) &= \frac{A_0^{(\alpha)}}{\Gamma(\bar{\beta}_1 - \bar{\beta}_{\alpha+1}) \left(\prod_{p=1}^{\alpha-2} \Gamma(\bar{\beta}_{p+1} - \bar{\beta}_{\alpha+p}) \right)} \\ &\times \sum_{n_1 n_2 \dots n_{\alpha-2}} \frac{\prod_{p=1}^{\alpha-2} [\Gamma(\bar{\beta}_{p+1} - \bar{\beta}_{\alpha+p} + n_p) \Gamma(\xi_{\alpha,p}(\mathbf{n}))]}{\left(\prod_{p=1}^{\alpha-3} \Gamma(\eta_{\alpha,p}(\mathbf{n})) \right) \Gamma(\zeta_{\alpha}(\mathbf{n})) \left(\prod_{p=1}^{\alpha-2} n_p! \right)} \\ &\times (1 - y)^{\zeta_{\alpha}(\mathbf{n}) - 1} {}_2F_1\left(\bar{\beta}_{\alpha} - \bar{\beta}_{2\alpha-1}, \eta_{\alpha,\alpha-2}(\mathbf{n}); \zeta_{\alpha}(\mathbf{n}); 1 - y\right), \end{aligned} \quad (4.19)$$

where

$$\begin{aligned} \xi_{\alpha,p}(\mathbf{n}) &= \sum_{q=1}^p (\bar{\beta}_q - \bar{\beta}_{\alpha+q} + n_q), & \eta_{\alpha,p}(\mathbf{n}) &= \sum_{q=1}^{p+1} (\bar{\beta}_q - \bar{\beta}_{\alpha+q}) + \sum_{q=1}^p n_q, \\ \zeta_{\alpha}(\mathbf{n}) &= \sum_{p=1}^{\alpha} \bar{\beta}_p - \sum_{p=1}^{\alpha-1} \bar{\beta}_{\alpha+p} + \sum_{p=1}^{\alpha-2} n_p - 1, \end{aligned} \quad (4.20)$$

if there exists some permutation $(i_1, i_2, \dots, i_\alpha)$ of $(1, 2, \dots, \alpha)$ such that

$$\bar{\beta}_{i_1} > \bar{\beta}_{\alpha+1}, \quad \bar{\beta}_{i_2} > \bar{\beta}_{\alpha+2}, \quad \dots, \quad \bar{\beta}_{i_{\alpha-1}} > \bar{\beta}_{2\alpha-1}, \quad \bar{\beta}_{i_\alpha} > 1. \quad (4.21)$$

The other weight functions $h_\mu^{(\alpha)}(y)$, $\mu = 1, 2, \dots, (\lambda - 2)/2$, and their respective conditions of applicability can be found from (4.19), (4.20), and (4.21) by making the substitutions

$$\begin{aligned} \bar{\beta}_1 &\rightarrow \bar{\beta}_{\mu+1}, & \bar{\beta}_2 &\rightarrow \bar{\beta}_{\mu+2}, & \dots, & \bar{\beta}_{\lambda-\mu-1} &\rightarrow \bar{\beta}_{\lambda-1}, & \bar{\beta}_{\lambda-\mu} &\rightarrow \bar{\beta}_1 + 1, \\ \bar{\beta}_{\lambda-\mu+1} &\rightarrow \bar{\beta}_2 + 1, & \dots, & \bar{\beta}_{\lambda-1} &\rightarrow \bar{\beta}_{\mu+1}. \end{aligned} \quad (4.22)$$

Setting $\alpha = 2$ in (4.19) and (4.20) gives back (4.18) as it should be. For $\alpha = 3$, Eq. (4.19) contains a single summation, which can be performed by using Eq. (I.33) of [46]. This leads to a closed expression of $h_0^{(3)}(y)$ in terms of an Appell function $F_3(\alpha, \alpha'; \beta, \beta'; \gamma; x, y)$,

$$\begin{aligned} h_0^{(3)}(y) &= A_0^{(3)} \frac{y^{\bar{\beta}_5 - \bar{\beta}_3} (1 - y)^{\bar{\beta}_1 + \bar{\beta}_2 + \bar{\beta}_3 - \bar{\beta}_4 - \bar{\beta}_5 - 2}}{\Gamma(\bar{\beta}_1 + \bar{\beta}_2 + \bar{\beta}_3 - \bar{\beta}_4 - \bar{\beta}_5 - 1)} F_3 \left(\bar{\beta}_1 - \bar{\beta}_4, \bar{\beta}_3 - \bar{\beta}_5; \bar{\beta}_2 - \bar{\beta}_4, \bar{\beta}_3 - 1; \right. \\ &\quad \left. \bar{\beta}_1 + \bar{\beta}_2 + \bar{\beta}_3 - \bar{\beta}_4 - \bar{\beta}_5 - 1; 1 - y, 1 - \frac{1}{y} \right). \end{aligned} \quad (4.23)$$

It should be noted that this result can also be obtained by direct integration without expanding the hypergeometric function into powers, as done in Appendix B, by using instead Eq. (2.21.1.20) of [47].

For higher α values, no closed expression of (4.19) has been proved to be valid in general, but we conjecture that it is given in terms of a Meijer G -function,

$$h_0^{(\alpha)}(y) = A_0^{(\alpha)} G_{\alpha, \alpha}^{\alpha, 0} \left(y \left| \begin{array}{c} \bar{\beta}_1 - 1, \dots, \bar{\beta}_\alpha - 1 \\ 0, \bar{\beta}_{\alpha+1} - 1, \dots, \bar{\beta}_{\lambda-1} - 1 \end{array} \right. \right). \quad (4.24)$$

Equation (4.24) has been demonstrated only for $\alpha = 1, 2$, and 3 , for which its equivalence to (4.17), (4.18), and (4.23) results from Eqs. (8.4.2.3), (8.4.49.22), and (8.4.51.3) of [47], respectively.

By making the parameter substitution (4.22) in (4.24) and (4.21), we get for $h_\mu^{(\alpha)}(y)$ and the corresponding applicability conditions the same results (4.15) and (4.14) as previously obtained when $\alpha \leq [(\lambda - 1)]/2$. In the present case, of course, we have to set $\lambda = 2\alpha$ in (4.15) and (4.14) and to restrict the range of y to the interval $(0, 1)$. At this point, it is worth noting that Eq. (4.15) would straightforwardly result from an unquestioning inversion of (4.4), but that the convolution process adopted here is essential to derive the positivity conditions (4.14).

Having found a solution to the problem stated in (4.4), we may now ask whether this solution is unique. An answer is provided by the (sufficient) condition of Carleman [41]: if a solution exists and $S \equiv \sum_{k=1}^{\infty} \left(B_{\mu}^{(\alpha)}(k) \right)^{-1/(2k)}$ diverges (resp. converges), then the solution is unique (resp. non-unique solutions may exist). As in [34], the possible convergence of S can be tested by applying the logarithmic test [47]. We obtain

$$\lim_{k \rightarrow \infty} \frac{\ln \left[\left(B_{\mu}^{(\alpha)}(k) \right)^{-1/(2k)} \right]}{\ln k} = - \left(\frac{\lambda}{2} - \alpha \right). \quad (4.25)$$

Hence S converges or diverges according to whether $-\left(\frac{\lambda}{2} - \alpha\right) < -1$ or $-\left(\frac{\lambda}{2} - \alpha\right) > -1$. We conclude that except in those cases where λ is odd and $\alpha = \frac{\lambda-1}{2}$, or λ is even and $\alpha = \frac{\lambda}{2}$, other solutions than those found hereabove may exist. To construct them, we might employ methods similar to those used in [34], but we shall pursue this subject no further, leaving it for a separate publication.

As a consequence of the unity resolution relation (4.1), we can express any CS $|z; \mu; \alpha\rangle$, defined in $\mathcal{F}_{\mu}$, in terms of any set of CS $|z; \mu; \alpha'\rangle$ ($\alpha' = \alpha$ or $\alpha' \neq \alpha$), defined in the same,

$$|z; \mu; \alpha\rangle = \int d\rho_{\mu}^{(\alpha')}(z', z'^*) |z'; \mu; \alpha'\rangle \langle z'; \mu; \alpha' | z; \mu; \alpha\rangle. \quad (4.26)$$

The reproducing kernel $\langle z'; \mu; \alpha' | z; \mu; \alpha\rangle$ is given in (3.18). More generally, an arbitrary element $|\psi_{\mu}\rangle$ of $\mathcal{F}_{\mu}$ can be written in terms of any set of CS $|z; \mu; \alpha\rangle$ defined in $\mathcal{F}_{\mu}$,

$$|\psi_{\mu}\rangle = \int d\rho_{\mu}^{(\alpha)}(z, z^*) \tilde{\psi}_{\mu}^{(\alpha)}(z, z^*) |z; \mu; \alpha\rangle, \quad (4.27)$$

where

$$\tilde{\psi}_{\mu}^{(\alpha)}(z, z^*) = \langle z; \mu; \alpha | \psi_{\mu}\rangle = \sum_{k=0}^{\infty} c_k(z, z^*; \mu; \alpha) (z^*)^k \langle k\lambda + \mu | \psi_{\mu}\rangle, \quad (4.28)$$

and $c_k(z, z^*; \mu; \alpha)$ is defined in (3.10) and (3.12).

4.1.2 Unity Resolution in $\mathcal{F}$

In the Fock space $\mathcal{F}$, the number states $|k\lambda + \mu\rangle$, $k = 0, 1, \dots$, $\mu = 0, 1, \dots, \lambda - 1$, form a complete set and their resolution of unity is given by $\sum_{\mu=0}^{\lambda-1} \sum_{k=0}^{\infty} |k\lambda + \mu\rangle \langle k\lambda + \mu| = \sum_{\mu=0}^{\lambda-1} P_{\mu} = I$.

The only set of CS $|z; \mu; \alpha\rangle$, $\mu = 0, 1, \dots, \lambda - \alpha - 1$, defined in the whole space $\mathcal{F}$ corresponds to $\alpha = 0$. Since two such states belonging to different subspaces $\mathcal{F}_{\mu}$ and $\mathcal{F}_{\mu'}$

are orthogonal (see (3.18)), it directly follows from (4.1) that this set of CS gives rise to a resolution of unity in $\mathcal{F}$ with a set of positive measures $d\rho_\mu^{(0)}(z, z^*)$, $\mu = 0, 1, \dots, \lambda - 1$,

$$\sum_{\mu=0}^{\lambda-1} \int d\rho_\mu^{(0)}(z, z^*) |z; \mu; 0\rangle \langle z; \mu; 0| = I. \quad (4.29)$$

Taking (3.3) into account, we recover the result previously obtained for the annihilation-operator CS of the C_λ -extended oscillator SGA [33].

An arbitrary element $|\psi\rangle = \sum_{\mu=0}^{\lambda-1} |\psi_\mu\rangle$ of $\mathcal{F}$, where $|\psi_\mu\rangle \in \mathcal{F}_\mu$, $\mu = 0, 1, \dots, \lambda - 1$, can be expanded in terms of the CS $|z; \mu; 0\rangle$, $\mu = 0, 1, \dots, \lambda - 1$,

$$|\psi\rangle = \sum_{\mu=0}^{\lambda-1} \int d\rho_\mu^{(0)}(z, z^*) \tilde{\psi}_\mu^{(0)}(z, z^*) |z; \mu; 0\rangle, \quad (4.30)$$

where $\tilde{\psi}_\mu^{(0)}(z, z^*)$ is defined in (4.28).

4.2 Unity Resolution Relation for the Coherent States $|z\rangle$

As mentioned at the end of Subsection 3.2, for values of λ greater than two, the eigenstates $|z\rangle$ of the C_λ -extended oscillator annihilation operator a may be considered as a generalization of paraboson CS. For the latter, Sharma *et al.* [35] demonstrated that a diagonal coherent state resolution of unity, similar to (4.1), does not exist, but found instead a nondiagonal one.

In the present Section, we will extend their approach to the case of the C_λ -extended oscillator CS and prove that they satisfy a unity resolution relation, which may be written in either form

$$\sum_{\mu=0}^{\lambda-1} \int d\rho_\mu(z, z^*) |z_\mu\rangle \langle z_\mu| = I \quad (4.31)$$

or

$$\sum_{\mu=0}^{\lambda-1} \int d\sigma_\mu(z, z^*) |z\rangle \langle z e^{2\pi i \mu / \lambda}| = I. \quad (4.32)$$

Here the integration is carried out over the whole complex plane,

$$d\rho_\mu(z, z^*) = \mathcal{N}(|z|) h_\mu(t) d^2 z, \quad d\sigma_\mu(z, z^*) = \mathcal{N}(|z|) g_\mu(t) d^2 z, \quad (4.33)$$

$$h_\mu(t) = \lambda^\lambda \left(\prod_{\nu=1}^{\mu} \bar{\beta}_\nu \right) t^{\lambda-\mu-1} h_\mu^{(0)}(t^\lambda), \quad (4.34)$$

$$g_\mu(t) = \frac{1}{\lambda} \sum_{\nu=0}^{\lambda-1} e^{2\pi i \mu \nu / \lambda} h_\nu(t), \quad (4.35)$$

$$|z_\mu\rangle = \left(\frac{N_\mu^{(0)}(|z|^\lambda)}{\mathcal{N}(|z|)} \right)^{1/2} \left(\prod_{\nu=1}^{\mu} \bar{\beta}_\nu \right)^{-1/2} \left(\frac{z}{\sqrt{\lambda}} \right)^\mu |z^\lambda; \mu; 0\rangle, \quad (4.36)$$

where $t = |z|^2/\lambda$, $d^2z = |z|d|z|d\phi$, and $h_\mu^{(0)}(y)$ is defined in (4.15). It should be noted that

$$|z\rangle = \sum_{\mu=0}^{\lambda-1} |z_\mu\rangle \quad (4.37)$$

and that the states $|z_\mu\rangle \in \mathcal{F}_\mu$, $\mu = 0, 1, \dots, \lambda - 1$, are orthogonal to each other, but are not properly normalized.

To prove (4.31), let us start from (4.29), where z is replaced by $\omega = z^\lambda$. By taking (3.12) and (4.2) into account, it can be rewritten as

$$\sum_{\mu=0}^{\lambda-1} \int d^2\omega h_\mu^{(0)}(|\omega|^2/\lambda^\lambda) N_\mu^{(0)}(|\omega|) |\omega; \mu; 0\rangle \langle \omega; \mu; 0| = I, \quad (4.38)$$

where $d^2\omega = |\omega|d|\omega|d\varphi$. We now make the change of variable $\omega = z^\lambda$, $d^2\omega = \lambda^2|z|^{2\lambda-2}d^2z$, and also allow for the fact that ω covers the complex plane λ times when z covers it once. Equation (4.38) becomes

$$\lambda \sum_{\mu=0}^{\lambda-1} \int d^2z |z|^{2\lambda-2} h_\mu^{(0)}(|z|^{2\lambda}/\lambda^\lambda) N_\mu^{(0)}(|z|^\lambda) |z^\lambda; \mu; 0\rangle \langle z^\lambda; \mu; 0| = I, \quad (4.39)$$

which directly leads to (4.31) after expressing $|z^\lambda; \mu; 0\rangle$ and $\langle z^\lambda; \mu; 0|$ in terms of $|z_\mu\rangle$ and $\langle z_\mu|$, respectively, and taking definitions (4.33) and (4.34) into account.

To transform (4.31) into (4.32), we note that from (4.36) and (4.37), it follows that

$$|ze^{2\pi i\mu/\lambda}\rangle = \sum_{\nu=0}^{\lambda-1} e^{2\pi i\mu\nu/\lambda} |z_\nu\rangle. \quad (4.40)$$

Inverting this equation leads to

$$|z_\nu\rangle = \frac{1}{\lambda} \sum_{\mu=0}^{\lambda-1} e^{-2\pi i\mu\nu/\lambda} |ze^{2\pi i\mu/\lambda}\rangle. \quad (4.41)$$

By inserting (4.41) and its Hermitian conjugate into (4.31), we obtain

$$\frac{1}{\lambda^2} \sum_{\mu, \mu', \nu=0}^{\lambda-1} e^{2\pi i(\mu' - \mu)\nu/\lambda} \int d^2z \mathcal{N}(|z|) h_\nu(t) |ze^{2\pi i\mu/\lambda}\rangle \langle ze^{2\pi i\mu'/\lambda}| = I. \quad (4.42)$$

It is now straightforward to get (4.32) by changing z into $z'e^{-2\pi i\mu/\lambda}$, $\mu' - \mu$ into $\mu'' \bmod \lambda$, and by performing the summation over μ .

We have therefore shown that the diagonal coherent state resolution of unity valid for the set of CS $\{|z; \mu; 0\rangle | \mu = 0, 1, \dots, \lambda - 1\}$ is entirely equivalent to a nondiagonal one for the CS

$|z\rangle$. Via a change of variable $z \rightarrow \omega$ and a redefinition of the component states $|z^\lambda; \mu; 0\rangle \rightarrow |z_\mu\rangle \rightarrow |ze^{2\pi i\mu/\lambda}\rangle$, the set of positive weight functions $\{h_\mu^{(0)}(y)|\mu = 0, 1, \dots, \lambda - 1\}$, has been converted into a set of linear combinations $\{g_\mu(t)|\mu = 0, 1, \dots, \lambda - 1\}$ with complex coefficients of well-behaved positive functions. Although this type of weight functions is unusual, the probabilistic interpretation of the weight is not lost provided one keeps the above-mentioned equivalence in mind.

As in Subsubsection 4.1.2, any vector of $\mathcal{F}$ can be expanded in terms of the CS $|z\rangle$,

$$|\psi\rangle = \sum_{\mu=0}^{\lambda-1} \int d\sigma_\mu(z, z^*) \tilde{\psi}\left(ze^{2\pi i\mu/\lambda}, z^*e^{-2\pi i\mu/\lambda}\right) |z\rangle, \quad (4.43)$$

where

$$\tilde{\psi}(z, z^*) = \langle z|\psi\rangle. \quad (4.44)$$

Equivalently, we may use (4.31) to express $|\psi\rangle$ as

$$|\psi\rangle = \sum_{\mu=0}^{\lambda-1} \int d\rho_\mu(z, z^*) \tilde{\psi}_\mu(z, z^*) |z_\mu\rangle, \quad (4.45)$$

where

$$\tilde{\psi}_\mu(z, z^*) = \langle z_\mu|\psi\rangle. \quad (4.46)$$

For $\lambda = 2$, the unity resolution relations (4.31) and (4.32) become

$$\int d\rho_0(z, z^*) |z_0\rangle \langle z_0| + \int d\rho_1(z, z^*) |z_1\rangle \langle z_1| = I \quad (4.47)$$

and

$$\int d\sigma_0(z, z^*) |z\rangle \langle z| + \int d\sigma_1(z, z^*) |z\rangle \langle -z| = I, \quad (4.48)$$

where

$$|z_0\rangle = \frac{1}{2}(|z\rangle + |-z\rangle), \quad |z_1\rangle = \frac{1}{2}(|z\rangle - |-z\rangle). \quad (4.49)$$

In such a case, $\mathcal{N}(|z|)$, $h_\mu(t)$, and $g_\mu(t)$ can be rewritten in terms of modified Bessel functions I_ν and K_ν as

$$\mathcal{N}(|z|) = 2^{\bar{\beta}_1-1} \Gamma(\bar{\beta}_1) |z|^{2(1-\bar{\beta}_1)} \left[I_{\bar{\beta}_1-1}(|z|^2) + I_{\bar{\beta}_1}(|z|^2) \right], \quad (4.50)$$

$$h_0(t) = 4t h_0^{(0)}(t^2) = \left[2^{\bar{\beta}_1-1} \pi \Gamma(\bar{\beta}_1) \right]^{-1} |z|^{2\bar{\beta}_1} K_{\bar{\beta}_1-1}(|z|^2), \quad (4.51)$$

$$h_1(t) = 4\bar{\beta}_1 h_1^{(0)}(t^2) = \left[2^{\bar{\beta}_1-1} \pi \Gamma(\bar{\beta}_1) \right]^{-1} |z|^{2\bar{\beta}_1} K_{\bar{\beta}_1}(|z|^2), \quad (4.52)$$

$$g_0(t) = \frac{1}{2}[h_0(t) + h_1(t)] = \left[2^{\bar{\beta}_1} \pi \Gamma(\bar{\beta}_1) \right]^{-1} |z|^{2\bar{\beta}_1} \left[K_{\bar{\beta}_1-1}(|z|^2) + K_{\bar{\beta}_1}(|z|^2) \right], \quad (4.53)$$

$$g_1(t) = \frac{1}{2}[h_0(t) - h_1(t)] = \left[2^{\bar{\beta}_1} \pi \Gamma(\bar{\beta}_1) \right]^{-1} |z|^{2\bar{\beta}_1} \left[K_{\bar{\beta}_1-1}(|z|^2) - K_{\bar{\beta}_1}(|z|^2) \right], \quad (4.54)$$

thus giving for $d\rho_\mu(z, z^*)$ and $d\sigma_\mu(z, z^*)$ the results previously obtained for paraboson CS [35],

$$d\rho_\mu(z, z^*) = \frac{|z|^2}{\pi} \left[I_{\bar{\beta}_1-1}(|z|^2) + I_{\bar{\beta}_1}(|z|^2) \right] K_{\bar{\beta}_1-1+\mu}(|z|^2) d^2z, \quad (4.55)$$

$$d\sigma_\mu(z, z^*) = \frac{|z|^2}{2\pi} \left[I_{\bar{\beta}_1-1}(|z|^2) + I_{\bar{\beta}_1}(|z|^2) \right] \left[K_{\bar{\beta}_1-1}(|z|^2) + (-1)^\mu K_{\bar{\beta}_1}(|z|^2) \right] d^2z. \quad (4.56)$$

5 BARGMANN REPRESENTATIONS

It is well known that for the harmonic oscillator, it is possible to find a realization of the Fock space wherein any vector is described by an entire function: this is the so-called Bargmann representation [38] associated with the harmonic oscillator CS. Similar realizations of other Lie algebras in Hilbert spaces of entire analytic functions have also been found in connection with various types of CS [9, 10]. Here we shall construct such realizations for the CS considered in the previous Sections.

5.1 Bargmann Representations Corresponding to the Coherent States $|z; \mu; \alpha\rangle$

5.1.1 Bargmann Representations of $\mathcal{F}_\mu$

As reviewed in Sec. 2, any subspace $\mathcal{F}_\mu$ of $\mathcal{F}$ carries a unirrep of the C_λ -extended oscillator SGA, characterized by the eigenvalue c_μ of C and by the lowest eigenvalue $(\mu + \gamma_\mu + \frac{1}{2})/\lambda = (\bar{\beta}_\mu + \bar{\beta}_{\mu+1})/2$ of J_0 . Furthermore, in Subsubsec. 4.1.1, it was proved that for any $\alpha \leq \lambda - \mu - 1$, the states $|z; \mu; \alpha\rangle$ form a complete (in fact, an overcomplete) set in $\mathcal{F}_\mu$ for some appropriate choice of the algebra parameters if z runs over the complex plane for $\alpha \leq [(\lambda - 1)/2]$ or over the unit disc for $\alpha = [\lambda/2]$ and λ even. We shall now proceed to show that with any such set, parametrized by a given α , we can associate a realization of $\mathcal{F}_\mu$ as a space $\mathcal{B}_\mu^{(\alpha)}$ of entire analytic functions, wherein the generators of the C_λ -extended oscillator SGA become differential operators.

For such a purpose, it is convenient to replace the normalized CS $|z; \mu; \alpha\rangle$ by unnormalized ones (denoted by a round bracket instead of an angular one),

$$|z; \mu; \alpha) \equiv \left[N_\mu^{(\alpha)}(|z|) \right]^{1/2} |z; \mu; \alpha\rangle, \quad (5.1)$$

and to substitute z^* for z . Then any vector $|\psi_\mu\rangle \in \mathcal{F}_\mu$ can be realized by the entire function

$$\begin{aligned}\psi_\mu^{(\alpha)}(z) &= (z^*; \mu; \alpha | \psi\rangle \\ &= \sum_{k=0}^{\infty} \left(\frac{\prod_{\nu=\mu+1}^{\mu+\alpha} (\bar{\beta}_\nu)_k}{k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_\nu + 1)_k \right) \left(\prod_{\nu=\mu+\alpha+1}^{\lambda-1} (\bar{\beta}_\nu)_k \right)} \right)^{1/2} \\ &\quad \times \langle k\lambda + \mu | \psi_\mu \rangle \left(\lambda^{-(\lambda-2\alpha)/2} z \right)^k,\end{aligned}\tag{5.2}$$

related to $\tilde{\psi}_\mu^{(\alpha)}(z, z^*)$, defined in (4.28), by

$$\tilde{\psi}_\mu^{(\alpha)}(z, z^*) = \left[N_\mu^{(\alpha)}(|z|) \right]^{-1/2} \psi_\mu^{(\alpha)}(z^*).\tag{5.3}$$

The functions (5.2) are the elements of a Hilbert space $\mathcal{B}_\mu^{(\alpha)}$. Its scalar product is defined by

$$\left(\psi_\mu'^{(\alpha)}, \psi_\mu^{(\alpha)} \right) = \int d^2z h_\mu^{(\alpha)}(y) \left[\psi_\mu'^{(\alpha)}(z) \right]^* \psi_\mu^{(\alpha)}(z), \quad y = |z|^2 / \lambda^{\lambda-2\alpha},\tag{5.4}$$

where the integral is extended over the complex plane or the unit disc according to the value of α . For any function $\psi_\mu^{(\alpha)}(z) \in \mathcal{B}_\mu^{(\alpha)}$, the condition $\left(\psi_\mu^{(\alpha)}, \psi_\mu^{(\alpha)} \right) < \infty$ is satisfied.

The functions

$$\begin{aligned}\varphi_{\mu,k}^{(\alpha)}(z) &= (z^*; \mu; \alpha | k\lambda + \mu\rangle \\ &= \left(\frac{\prod_{\nu=\mu+1}^{\mu+\alpha} (\bar{\beta}_\nu)_k}{k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_\nu + 1)_k \right) \left(\prod_{\nu=\mu+\alpha+1}^{\lambda-1} (\bar{\beta}_\nu)_k \right)} \right)^{1/2} \left(\lambda^{-(\lambda-2\alpha)/2} z \right)^k,\end{aligned}\tag{5.5}$$

representing the number-state vectors $|k\lambda + \mu\rangle$, $k = 0, 1, \dots$, form an orthonormal basis of $\mathcal{B}_\mu^{(\alpha)}$. Equation (5.2) can therefore be rewritten as

$$\psi_\mu^{(\alpha)}(z) = \sum_{k=0}^{\infty} \langle k\lambda + \mu | \psi_\mu \rangle \varphi_{\mu,k}^{(\alpha)}(z).\tag{5.6}$$

Any operator X defined in $\mathcal{F}_\mu$ is represented in $\mathcal{B}_\mu^{(\alpha)}$ by some differential operator $\mathcal{X}_\mu^{(\alpha)}$, defined by

$$(z^*; \mu; \alpha | X | \psi_\mu \rangle = \mathcal{X}_\mu^{(\alpha)} \psi_\mu^{(\alpha)}(z)\tag{5.7}$$

for any $|\psi_\mu\rangle \in \mathcal{F}_\mu$. For the number operator N , we get

$$\mathcal{N}_\mu^{(\alpha)} = \lambda z \frac{\partial}{\partial z} + \mu.\tag{5.8}$$

Similarly, for the generators J_+ , J_- , and J_0 of the C_λ -extended oscillator SGA, we obtain

$$\mathcal{J}_{+\mu}^{(\alpha)} = \lambda^{\alpha-1} z \prod_{\nu=\mu+1}^{\mu+\alpha} \left(z \frac{\partial}{\partial z} + \bar{\beta}_\nu \right), \quad (5.9)$$

$$\mathcal{J}_{-\mu}^{(\alpha)} = \lambda^{\lambda-\alpha-1} \left[\prod_{\nu=1}^{\mu} \left(z \frac{\partial}{\partial z} + \bar{\beta}_\nu + 1 \right) \right] \left[\prod_{\nu=\mu+\alpha+1}^{\lambda-1} \left(z \frac{\partial}{\partial z} + \bar{\beta}_\nu \right) \right] \frac{\partial}{\partial z}, \quad (5.10)$$

$$\mathcal{J}_{0\mu}^{(\alpha)} = z \frac{\partial}{\partial z} + \frac{1}{2}(\bar{\beta}_\mu + \bar{\beta}_{\mu+1}). \quad (5.11)$$

In deriving these equations, use has been made of the expansion (5.6) and of the known action of N , $(a^\dagger)^\lambda$, a^λ , and H_0 on $|k\lambda + \mu\rangle$ (see (2.10), (3.9), (3.8) and (2.11), respectively). It is straightforward to check that the differential operators (5.9)–(5.11) fulfil the commutation relations (2.15) characterizing the C_λ -extended oscillator SGA. In Appendix C, it is proved that they also satisfy the Hermiticity properties (2.14) with respect to the scalar product (5.4) of $\mathcal{B}_\mu^{(\alpha)}$.

It should be noted that in the $\alpha = 0$ case, $\mathcal{J}_{+\mu}^{(0)}$ is essentially the operator that acts on $\mathcal{B}_\mu^{(0)}$ as a multiplication by z , whereas $\mathcal{J}_{-\mu}^{(0)}$ is a λ th-order differential operator. If in addition $\lambda = 2$, we get back the well-known realization of $\mathfrak{su}(1,1)$ connected with the Barut-Girardello CS [5], corresponding to the unirrep characterized by $\frac{1}{2}(\bar{\beta}_\mu + \bar{\beta}_{\mu+1}) = \frac{1}{2}(\bar{\beta}_1 + \mu)$ and c_μ ,

$$\mathcal{J}_{+\mu}^{(0)} = \frac{1}{2}z, \quad \mathcal{J}_{-\mu}^{(0)} = 2 \left(z \frac{\partial}{\partial z} + \bar{\beta}_1 + \mu \right) \frac{\partial}{\partial z}, \quad \mathcal{J}_{0\mu}^{(0)} = z \frac{\partial}{\partial z} + \frac{1}{2}(\bar{\beta}_1 + \mu). \quad (5.12)$$

Similarly, for $\lambda = 2$, $\alpha = 1$, and $\mu = 0$, $\mathcal{J}_{-0}^{(1)}$ is the operator that acts on $\mathcal{B}_0^{(1)}$ as a derivation $\partial/\partial z$. This corresponds to the realization of $\mathfrak{su}(1,1)$ connected with the Perelomov CS [9], corresponding to the unirrep specified by $\frac{1}{2}\bar{\beta}_1$ and c_0 ,

$$\mathcal{J}_{+0}^{(1)} = z \left(z \frac{\partial}{\partial z} + \bar{\beta}_1 \right), \quad \mathcal{J}_{-0}^{(1)} = \frac{\partial}{\partial z}, \quad \mathcal{J}_{00}^{(1)} = z \frac{\partial}{\partial z} + \frac{1}{2}\bar{\beta}_1. \quad (5.13)$$

5.1.2 Bargmann Representation of $\mathcal{F}$

For $\alpha = 0$, we may realize the whole Fock space $\mathcal{F} = \sum_{\mu=0}^{\lambda-1} \oplus \mathcal{F}_\mu$ as a Hilbert space $\mathcal{B}^{(0)} = \sum_{\mu=0}^{\lambda-1} \oplus \mathcal{B}_\mu^{(0)}$ of entire analytic functions. It is then convenient to introduce vector CS [10, 39], defined as row vectors

$$||z; 0\rangle\rangle = (|z; 0; 0\rangle, |z; 1; 0\rangle, \dots, |z; \lambda - 1; 0\rangle), \quad (5.14)$$

the corresponding bras being column vectors

$$((z; 0|) = \begin{pmatrix} (z; 0; 0| \\ (z; 1; 0| \\ \vdots \\ (z; \lambda - 1; 0| \end{pmatrix}. \quad (5.15)$$

The unity resolution relation in $\mathcal{F}$, given in (4.29), can be rewritten as

$$\int d^2 z ||z; 0)) h^{(0)}(y) ((z; 0| = I, \quad (5.16)$$

where $h^{(0)}(y) = \text{diag} (h_0^{(0)}(y), h_1^{(0)}(y), \dots, h_{\lambda-1}^{(0)}(y))$ is a diagonal $\lambda \times \lambda$ matrix.

Any vector $|\psi\rangle = \sum_{\mu=0}^{\lambda-1} |\psi_\mu\rangle \in \mathcal{F}$ is realized by a column of entire functions

$$\psi^{(0)}(z) = ((z^*; 0| |\psi\rangle = \begin{pmatrix} \psi_0^{(0)}(z) \\ \psi_1^{(0)}(z) \\ \vdots \\ \psi_{\lambda-1}^{(0)}(z) \end{pmatrix}, \quad (5.17)$$

and the scalar product in $\mathcal{B}^{(0)}$ reads

$$(\psi'^{(0)}, \psi^{(0)}) = \int d^2 z [\psi'^{(0)}(z)]^\dagger h^{(0)}(y) \psi^{(0)}(y). \quad (5.18)$$

Any operator X defined in $\mathcal{F}$ is represented in $\mathcal{B}^{(0)}$ by an operator-valued $\lambda \times \lambda$ matrix $\mathcal{X}^{(0)}$, defined by

$$((z^*; 0| |X| \psi\rangle = \mathcal{X}^{(0)} \psi^{(0)}(z). \quad (5.19)$$

If X does not produce transitions between different subspaces $\mathcal{F}_\mu$, then $\mathcal{X}^{(0)}$ is a diagonal matrix. Such is the case for the projection operators P_μ , which are represented by

$$\mathcal{P}_\mu = e_{\mu+1, \mu+1}, \quad (5.20)$$

where $e_{\mu\nu}$ denotes the matrix with entry one in row μ and column ν , and zeros everywhere else. The diagonal matrix $\mathcal{X}^{(0)}$ may then be written as $\mathcal{X}^{(0)} = \sum_\mu \mathcal{X}_\mu^{(0)} \mathcal{P}_\mu$, where $\mathcal{X}_\mu^{(0)}$ is the operator representing in $\mathcal{B}_\mu^{(0)}$ the restriction of X to $\mathcal{F}_\mu$. In particular, for the generators of the C_λ -extended oscillator SGA, we find

$$\mathcal{J}_q^{(0)} = \sum_\mu \mathcal{J}_{q\mu}^{(0)} \mathcal{P}_\mu, \quad q = +, -, 0, \quad (5.21)$$

where $\mathcal{J}_{q\mu}^{(0)}$, $q = +, -, 0$, are given in (5.9)–(5.11). Similarly, the number operator N is represented by

$$\mathcal{N}^{(0)} = \sum_{\mu} \mathcal{N}_{\mu}^{(0)} \mathcal{P}_{\mu}, \quad (5.22)$$

where $\mathcal{N}_{\mu}^{(0)}$ is defined in (5.8).

On the contrary, operators giving rise to transitions between different subspaces $\mathcal{F}_{\mu}$ are represented by nondiagonal operator-valued matrices. For the creation and annihilation operators of the C_{λ} -extended oscillator, for instance, we obtain the matrices

$$\mathcal{A}^{\dagger(0)} = \begin{pmatrix} 0 & 0 & \dots & 0 & z/\sqrt{\lambda^{\lambda-1}\bar{\beta}_1\bar{\beta}_2\dots\bar{\beta}_{\lambda-1}} \\ \sqrt{\lambda\bar{\beta}_1} & 0 & \dots & 0 & 0 \\ 0 & \sqrt{\lambda\bar{\beta}_2} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & \sqrt{\lambda\bar{\beta}_{\lambda-1}} & 0 \end{pmatrix}, \quad (5.23)$$

$$\mathcal{A}^{(0)} = \begin{pmatrix} 0 & \sqrt{\lambda/\bar{\beta}_1}(z\frac{\partial}{\partial z} + \bar{\beta}_1) & 0 & \dots & 0 \\ 0 & 0 & \sqrt{\lambda/\bar{\beta}_2}(z\frac{\partial}{\partial z} + \bar{\beta}_2) & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \sqrt{\lambda/\bar{\beta}_{\lambda-1}}(z\frac{\partial}{\partial z} + \bar{\beta}_{\lambda-1}) \\ \sqrt{\lambda^{\lambda+1}\bar{\beta}_1\bar{\beta}_2\dots\bar{\beta}_{\lambda-1}}\frac{\partial}{\partial z} & 0 & 0 & \dots & 0 \end{pmatrix}, \quad (5.24)$$

respectively. It is straightforward to check that the operator-valued matrices (5.20) and (5.22)–(5.24) satisfy the commutation relations (2.3) of the C_{λ} -extended oscillator algebra. The more difficult check of their Hermiticity properties with respect to the scalar product (5.18) of $\mathcal{B}^{(0)}$ is reviewed in Appendix C.

From (5.22)–(5.24), it is easy to recover (5.21) by using the relations

$$\mathcal{J}_+^{(0)} = \frac{1}{\lambda} (\mathcal{A}^{\dagger(0)})^{\lambda}, \quad \mathcal{J}_-^{(0)} = \frac{1}{\lambda} (\mathcal{A}^{(0)})^{\lambda}, \quad \mathcal{J}_0^{(0)} = \frac{1}{\lambda} \left[\mathcal{N}^{(0)} + \frac{1}{2} \sum_{\mu} (\beta_{\mu} + \beta_{\mu+1} + 1) \mathcal{P}_{\mu} \right], \quad (5.25)$$

corresponding to (2.12).

5.2 Bargmann Representation Corresponding to the Coherent States $|z\rangle$

In $\mathcal{F}$, we may alternatively use the unnormalized CS

$$|z\rangle \equiv [\mathcal{N}(|z|)]^{1/2}|z\rangle = \sum_{\mu} |z_{\mu}\rangle, \quad |z_{\mu}\rangle \equiv [\mathcal{N}(|z|)]^{1/2}|z_{\mu}\rangle. \quad (5.26)$$

Employing a vector notation again, let us define

$$||z\rangle\rangle = (|z_0\rangle, |z_1\rangle, \dots, |z_{\lambda-1}\rangle), \quad ((z|| = \begin{pmatrix} \langle z_0| \\ \langle z_1| \\ \vdots \\ \langle z_{\lambda-1}| \end{pmatrix}, \quad (5.27)$$

which are related to the row and column vectors (5.14) and (5.15) through the relations

$$||z\rangle\rangle = ||\omega; 0\rangle\rangle D(z), \quad ((z|| = D(z^*)((\omega; 0||. \quad (5.28)$$

Here $D(z)$ is a $\lambda \times \lambda$ diagonal matrix, given by

$$\begin{aligned} D(z) &= \text{diag}(D_0(z), D_1(z), \dots, D_{\lambda-1}(z)), \\ D_{\mu}(z) &= \left(\prod_{\nu=1}^{\mu} \bar{\beta}_{\nu} \right)^{-1/2} \left(\frac{z}{\sqrt{\lambda}} \right)^{\mu}, \quad \mu = 0, 1, \dots, \lambda - 1. \end{aligned} \quad (5.29)$$

The unity resolution relation in $\mathcal{F}$, given in (4.31), can be rewritten as

$$\int d^2 z ||z\rangle\rangle h(t) ((z|| = I, \quad (5.30)$$

where $h(t) = \text{diag}(h_0(t), h_1(t), \dots, h_{\lambda-1}(t))$.

By using the new vector CS (5.27), we get a new realization of $\mathcal{F}$ as a space $\mathcal{B}$ of entire analytic functions. Any vector $|\psi\rangle \in \mathcal{F}$ is realized by a column of entire functions

$$\psi(z) = ((z^*||\psi\rangle = D(z)\psi^{(0)}(\omega), \quad (5.31)$$

where $\psi^{(0)}(\omega)$ is defined in (5.17). The scalar product in $\mathcal{B}$ reads

$$(\psi', \psi) = \int d^2 z [\psi'(z)]^{\dagger} h(t) \psi(z). \quad (5.32)$$

Any operator X defined in $\mathcal{F}$ is represented in $\mathcal{B}$ by an operator-valued matrix $\mathcal{X}$, defined by

$$((z^*||X|\psi\rangle = \mathcal{X}\psi(z), \quad (5.33)$$

and given by

$$\mathcal{X} = D(z)\mathcal{X}^{(0)}D^{-1}(z) \quad (5.34)$$

in terms of $\mathcal{X}^{(0)}$ of Eq. (5.19).

From (5.9)–(5.11) and (5.21)–(5.24), it is straightforward to derive the following results:

$$\mathcal{J}_q = \sum_{\mu} \mathcal{J}_{q\mu} \mathcal{P}_{\mu}, \quad q = +, -, 0, \quad (5.35)$$

$$\mathcal{J}_{+\mu} = \frac{1}{\lambda} z^{\lambda}, \quad (5.36)$$

$$\mathcal{J}_{-\mu} = \frac{1}{\lambda} \left[\prod_{\nu=\mu+1}^{\lambda-1} \left(\frac{\partial}{\partial z} + \frac{\beta_{\nu}}{z} \right) \right] \frac{\partial}{\partial z} \left[\prod_{\nu=1}^{\mu} \left(\frac{\partial}{\partial z} + \frac{\beta_{\nu}}{z} \right) \right], \quad (5.37)$$

$$\mathcal{J}_{0\mu} = \frac{1}{\lambda} \left(z \frac{\partial}{\partial z} + \frac{1}{2} (\beta_{\mu} + \beta_{\mu+1} + 1) \right), \quad (5.38)$$

$$\mathcal{N} = \sum_{\mu} \mathcal{N}_{\mu} \mathcal{P}_{\mu}, \quad \mathcal{N}_{\mu} = z \frac{\partial}{\partial z}, \quad (5.39)$$

$$\mathcal{A}^{\dagger} = \begin{pmatrix} 0 & 0 & \dots & 0 & z \\ z & 0 & \dots & 0 & 0 \\ 0 & z & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & z & 0 \end{pmatrix}, \quad (5.40)$$

$$\mathcal{A} = \begin{pmatrix} 0 & \frac{\partial}{\partial z} + \frac{\beta_1}{z} & 0 & \dots & 0 \\ 0 & 0 & \frac{\partial}{\partial z} + \frac{\beta_2}{z} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \frac{\partial}{\partial z} + \frac{\beta_{\lambda-1}}{z} \\ \frac{\partial}{\partial z} & 0 & 0 & \dots & 0 \end{pmatrix}. \quad (5.41)$$

In particular, for $\lambda = 2$, we get back known results for paraboson CS [35]:

$$\mathcal{N} = \begin{pmatrix} z \frac{\partial}{\partial z} & 0 \\ 0 & z \frac{\partial}{\partial z} \end{pmatrix}, \quad \mathcal{A}^{\dagger} = \begin{pmatrix} 0 & z \\ z & 0 \end{pmatrix}, \quad \mathcal{A} = \begin{pmatrix} 0 & \frac{\partial}{\partial z} + \frac{\beta_1}{z} \\ \frac{\partial}{\partial z} & 0 \end{pmatrix}. \quad (5.42)$$

6 PHYSICAL APPLICATIONS

In this Section, we shall investigate the statistical and squeezing properties of the C_{λ} -extended oscillator CS $|z; \mu; \alpha\rangle$ and $|z\rangle$ with special emphasis on the comparison with those of the standard harmonic oscillator, which are retrieved for vanishing parameters α_{μ} .

6.1 Photon Statistics

For the conventional harmonic oscillator CS [1, 2, 3], the probability distribution of the photon number is Poissonian. Experimentally, however, the photon number statistics of real lasers is not exactly Poissonian [48]. Furthermore, various nonlinear interactions give rise to deviations from the Poisson distribution [11, 12]. A measure of such deviations is the Mandel parameter [49]

$$Q = \frac{(\Delta N_b)^2 - \langle N_b \rangle}{\langle N_b \rangle}, \quad (\Delta N_b)^2 \equiv \langle N_b^2 \rangle - \langle N_b \rangle^2, \quad (6.1)$$

which vanishes for the Poisson distribution, is positive for a super-Poissonian distribution (bunching effect), and negative for a sub-Poissonian distribution (antibunching effect). As mentioned in Sec. 1, in the present case we have $N_b = N$.

6.1.1 Statistical Properties of the Coherent States $|z; \mu; \alpha\rangle$

By using the Bargmann representation $\mathcal{N}_\mu^{(\alpha)}$ of N , given in (5.8), the averages of N and N^2 in the states $|z; \mu; \alpha\rangle$ can be expressed in terms of derivatives of the normalization coefficients $N_\mu^{(\alpha)}(|z|)$,

$$\begin{aligned} \langle z; \mu; \alpha | N^r | z; \mu; \alpha \rangle &= [N_\mu^{(\alpha)}(|z|)]^{-1} \left(\lambda z^* \frac{\partial}{\partial z^*} + \mu \right)^r N_\mu^{(\alpha)}(|z|) \\ &= [N_\mu^{(\alpha)}(|z|)]^{-1} \left(\lambda y \frac{\partial}{\partial y} + \mu \right)^r N_\mu^{(\alpha)}(|z|), \end{aligned} \quad (6.2)$$

where $r = 1, 2$, and $y = |z|^2 / \lambda^{\lambda-2\alpha}$.

For $\lambda = 2$, $\alpha = 1$ (and $\mu = 0$), i.e., for the Perelomov $\text{su}(1,1)$ CS given in (3.6), the normalization coefficient $N_0^{(1)}(|z|)$ takes a very simple form. The same is true for the Mandel parameter, which is independent of the parameter α_0 and is given by

$$Q = \frac{1+y}{1-y}, \quad (6.3)$$

where $y = |z|^2$. The distribution is then super-Poissonian as $1 < Q < \infty$ for $0 < y < 1$.

For the remaining CS $|z; \mu; \alpha\rangle$, the Mandel parameter can be written as

$$Q = \begin{cases} \lambda \left[1 - \bar{\beta}_{\lambda-1} - \left(\prod_{\nu=1}^{\alpha} \bar{\beta}_\nu \right) \left(\prod_{\nu=\alpha+1}^{\lambda-1} \bar{\beta}_\nu \right)^{-1} y \Phi_0^{\lambda-1}(y; 0; \alpha) + \bar{\beta}_{\lambda-1} \Phi_{\lambda-1}^{\lambda-2}(y; 0; \alpha) \right] \\ -1 & \text{if } \mu = 0, \end{cases}$$

$$\begin{aligned}
&= \left[\lambda^{-1} - \bar{\beta}_1 + \bar{\beta}_1 \Phi_1^0(y; 1; \alpha) \right]^{-1} \left\{ \left(\bar{\beta}_1 - \lambda^{-1} \right) \left[1 + \lambda \bar{\beta}_1 \Phi_1^0(y; 1; \alpha) \right] \right. \\
&\quad \left. - \lambda \bar{\beta}_1^2 \left[\Phi_1^0(y; 1; \alpha) \right]^2 + \lambda \left(\prod_{\nu=2}^{\alpha+1} \bar{\beta}_\nu \right) \left(\prod_{\nu=\alpha+2}^{\lambda-1} \bar{\beta}_\nu \right)^{-1} y \Phi_1^{\lambda-1}(y; 1; \alpha) \right\} \quad \text{if } \mu = 1, \\
&= \left[\mu \lambda^{-1} - \bar{\beta}_\mu + \bar{\beta}_\mu \Phi_\mu^{\mu-1}(y; \mu; \alpha) \right]^{-1} \left\{ \bar{\beta}_\mu - \mu \lambda^{-1} + \lambda \bar{\beta}_\mu \left(\bar{\beta}_\mu - \bar{\beta}_{\mu-1} - \lambda^{-1} \right) \Phi_\mu^{\mu-1}(y; \mu; \alpha) \right. \\
&\quad \left. - \lambda \bar{\beta}_\mu^2 \left[\Phi_\mu^{\mu-1}(y; \mu; \alpha) \right]^2 + \lambda \bar{\beta}_{\mu-1} \bar{\beta}_\mu \Phi_\mu^{\mu-2}(y; \mu; \alpha) \right\} \quad \text{if } \mu = 2, 3, \dots, \lambda - 1, \quad (6.4)
\end{aligned}$$

where

$$\Phi_\nu^{\nu'}(y; \mu; \alpha) = \frac{{}_\alpha F_{\lambda-\alpha-1} \left(\bar{\beta}_{\mu+1}^{(\nu')}, \dots, \bar{\beta}_{\mu+\alpha}^{(\nu')}; \bar{\beta}_1^{(\nu')}, \dots, \bar{\beta}_\mu^{(\nu')}, \bar{\beta}_{\mu+\alpha+1}^{(\nu')}, \dots, \bar{\beta}_{\lambda-1}^{(\nu')}; y \right)}{{}_\alpha F_{\lambda-\alpha-1} \left(\bar{\beta}_{\mu+1}^{(\nu)}, \dots, \bar{\beta}_{\mu+\alpha}^{(\nu)}; \bar{\beta}_1^{(\nu)}, \dots, \bar{\beta}_\mu^{(\nu)}, \bar{\beta}_{\mu+\alpha+1}^{(\nu)}, \dots, \bar{\beta}_{\lambda-1}^{(\nu)}; y \right)} \quad (6.5)$$

and

$$\begin{aligned}
\bar{\beta}_\mu^{(\nu)} &= \bar{\beta}_\mu + 1 & \text{if } \mu \leq \nu, \\
&= \bar{\beta}_\mu & \text{if } \mu > \nu.
\end{aligned} \quad (6.6)$$

For $\alpha = 0$, Eq. (6.5) reduces to

$$\Phi_\nu^{\nu'}(y; \mu; 0) = \frac{N_{\nu'}^{(0)}(|z|)}{N_\nu^{(0)}(|z|)}, \quad (6.7)$$

which is independent of μ . We then get back the results previously derived for the CS (3.3) of the C_λ -extended oscillator SGA (see Eq. (36) of [33]). For such states, it has been shown that the parameters α_μ have a very strong effect on Q . For instance, the super-Poissonian distribution obtained for $\mu = 0$ when $\alpha_\mu = 0$ (or $\bar{\beta}_\mu = \mu/\lambda$), $\mu = 0, 1, \dots, \lambda - 1$, is changed into a sub-Poissonian one when the α_μ 's are varied in an appropriate way, e.g., for $\alpha_0 < 0$ corresponding to $\bar{\beta}_1 < \lambda^{-1}$.

For $\alpha = 1$, let us consider the case where $\lambda = 3$, corresponding to the CS (3.13) and (3.14).

For $\mu = 0$, Q goes to 2 for both $y \rightarrow 0$ and $y \rightarrow \infty$, and any parameter values. Since the confluent hypergeometric function ${}_1F_1(a + n; a; y)$, $n \in \mathbb{N}$, reduces to an exponential multiplied by an n th-degree polynomial [45], it is easy to get simple analytical formulae for $\bar{\beta}_1 = \bar{\beta}_2 + n$ (or $\alpha_1 = -1 - 3n$). For instance

$$\begin{aligned}
Q &= 2 & \text{if } \bar{\beta}_1 = \bar{\beta}_2, \\
&= 2 - \frac{3y}{(y + \bar{\beta}_2)(y + \bar{\beta}_2 + 1)} & \text{if } \bar{\beta}_1 = \bar{\beta}_2 + 1.
\end{aligned} \quad (6.8)$$

In the latter case, $Q \leq 2$ for any value of $|z|$ and its minimum, $Q_{min} = 2 - 3 \left(\sqrt{\bar{\beta}_2} + \sqrt{\bar{\beta}_2 + 1} \right)^{-2}$, decreases from 2 to -1 when $\bar{\beta}_2$ decreases from $+\infty$ to 0. For other values of $\bar{\beta}_1$, a numerical study confirms this trend by showing that $Q \geq 2$ or $Q \leq 2$ on the whole range of $|z|$ according to whether $\bar{\beta}_1 < \bar{\beta}_2$ or $\bar{\beta}_1 > \bar{\beta}_2$. In particular, as shown in Fig. 3a, the strong super-Poissonian character of the distribution obtained for vanishing α_μ parameters (i.e., $\bar{\beta}_1 = 1/3$, $\bar{\beta}_2 = 2/3$) can be changed into a sub-Poissonian one over a limited range of $|z|$ values provided α_1 is negative enough (or $\bar{\beta}_2$ small enough).

For $\mu = 1$, Q goes to -1 or 2 for any parameter values according to whether $y \rightarrow 0$ or $y \rightarrow \infty$. Simple analytical formulae can be obtained for $\bar{\beta}_2 = \bar{\beta}_1 + 1 + n$ (or $\alpha_1 = 2 + 3n$), $n \in \mathbb{N}$. For instance

$$Q = 2 - \frac{3}{3y + 1} \quad \text{if } \bar{\beta}_2 = \bar{\beta}_1 + 1. \quad (6.9)$$

In such a case, Q is independent of $\bar{\beta}_1$ and increases from -1 to 2 . For other values of $\bar{\beta}_2$, a numerical study shows that if $\bar{\beta}_2 < \bar{\beta}_1 + 1$, Q has a maximum greater than 2, whereas if $\bar{\beta}_2 > \bar{\beta}_1 + 1$, its behaviour is rather similar to that of (6.9). Hence, as shown in Fig. 3b, large negative values of α_1 (or small values of $\bar{\beta}_2$) again drastically change Q as compared with the case of vanishing α_μ parameters.

6.1.2 Statistical Properties of the Coherent States $|z\rangle$

The averages of N and N^2 in the CS $|z\rangle$ are easily calculated from those determined in the previous Subsubsection since such states are linear combinations of the CS $|\omega; \mu; 0\rangle$, where $\omega = z^\lambda$. The result for Q reads

$$Q = \frac{S_2(|z|)}{S_1(|z|)} - \lambda \frac{S_1(|z|)}{\mathcal{N}(|z|)}, \quad (6.10)$$

where

$$S_1(|z|) = \sum_{\mu=0}^{\lambda-1} N_\mu^{(0)}(|\omega|) (t + \mu\lambda^{-1} - \bar{\beta}_\mu) \frac{t^\mu}{\prod_{\nu=1}^{\mu} \bar{\beta}_\nu}, \quad (6.11)$$

$$S_2(|z|) = \sum_{\mu=0}^{\lambda-1} N_\mu^{(0)}(|\omega|) \left[\mu(\mu-1)\lambda^{-1} - (2\mu-1)\bar{\beta}_\mu + \lambda\bar{\beta}_\mu^2 + (2\mu+1 - \lambda\bar{\beta}_\mu - \lambda\bar{\beta}_{\mu+1})t + \lambda t^2 \right] \frac{t^\mu}{\prod_{\nu=1}^{\mu} \bar{\beta}_\nu}, \quad (6.12)$$

and $t = |z|^2/\lambda$, $|\omega| = |z|^\lambda$.

For $\lambda = 2$, Q can be rewritten in terms of Bessel modified functions I_ν as

$$Q = (1 - 2\bar{\beta}_1)[2t - (1 - 2\bar{\beta}_1)R(t)]^{-1}[2t - 2(2t + \bar{\beta}_1)R(t) - (1 - 2\bar{\beta}_1)R^2(t)], \quad (6.13)$$

where

$$R(t) = \frac{I_{\bar{\beta}_1}(2t)}{I_{\bar{\beta}_1-1}(2t) + I_{\bar{\beta}_1}(2t)} \quad (t = |z|^2/2). \quad (6.14)$$

For $\bar{\beta}_1 = 1/2$ (or $\alpha_0 = 0$), we obtain $Q = 0$, as it should be, since the CS $|z\rangle$ then reduce to the conventional harmonic oscillator CS. In general, we find that Q behaves as $(2\bar{\beta}_1 - 1)|z|^2/(2\bar{\beta}_1)$ if $|z| \ll 1$, and as $(2\bar{\beta}_1 - 1)/(2|z|^2)$ if $|z| \gg 1$. As shown in Fig. 4a, the distribution is super-Poissonian or sub-Poissonian according to whether $\alpha_0 > 0$ (i.e., $\bar{\beta}_1 > 1/2$) or $\alpha_0 < 0$ (i.e., $\bar{\beta}_1 < 1/2$).

For higher values of λ , the behaviour of Q becomes more complicated as it now depends on more than one parameter. For $\lambda = 3$ and $|z| \ll 1$, for instance, Q behaves as $(2\bar{\beta}_1 - \bar{\beta}_2)|z|^2/(3\bar{\beta}_1\bar{\beta}_2)$ if $\bar{\beta}_2 \neq 2\bar{\beta}_1$, and as $(3\bar{\beta}_1 - 1)|z|^4/(18\bar{\beta}_1^2)$ if $\bar{\beta}_2 = 2\bar{\beta}_1 \neq 2/3$. Hence values of $\bar{\beta}_1$ smaller (resp. greater) than both $\bar{\beta}_2/2$ and $1/3$ favour sub-Poissonian (resp. super-Poissonian) distributions. In terms of the α_μ parameters, this means $\alpha_0 < 0$ and $\alpha_1 > \alpha_0$ (resp. $\alpha_0 > 0$ and $\alpha_1 < \alpha_0$). However, as shown in Fig. 4b, intermediate values of $\bar{\beta}_1$ may lead to an oscillating behaviour of Q .

6.2 Squeezing Properties

For conventional photon operators, the electromagnetic field components x_b and p_b are given by

$$x_b = \frac{1}{\sqrt{2}}(b + b^\dagger), \quad p_b = \frac{1}{i\sqrt{2}}(b - b^\dagger). \quad (6.15)$$

Their variances $(\Delta x_b)^2 = \langle x_b^2 \rangle - \langle x_b \rangle^2$ and $(\Delta p_b)^2 = \langle p_b^2 \rangle - \langle p_b \rangle^2$ in any state satisfy the conventional uncertainty relation

$$(\Delta x_b)^2(\Delta p_b)^2 \geq \frac{1}{4}, \quad (6.16)$$

the lower bound being attained by the vacuum state, for which $(\Delta x_b)_0^2 = (\Delta p_b)_0^2 = 1/2$. The quadrature x_b (resp. p_b) is said to be squeezed in a given state if $(\Delta x_b)^2 < (\Delta x_b)_0^2$ (resp. $(\Delta p_b)^2 < (\Delta p_b)_0^2$) or, in other words, if the ratio $X \equiv (\Delta x_b)^2/(\Delta x_b)_0^2$ (resp. $P \equiv (\Delta p_b)^2/(\Delta p_b)_0^2$) is less than one.

The C_λ -extended oscillator creation and annihilation operators $a^\dagger$, a may be regarded as describing “dressed” photons, which may have some applications in phenomenological models explaining non-intuitive observable phenomena [18]. It is therefore also interesting to consider deformed electromagnetic field components x and p , given by

$$x = \frac{1}{\sqrt{2}} (a + a^\dagger), \quad p = \frac{1}{i\sqrt{2}} (a - a^\dagger), \quad (6.17)$$

where

$$a = b \left(\frac{F(N_b)}{N_b} \right)^{1/2}, \quad a^\dagger = \left(\frac{F(N_b)}{N_b} \right)^{1/2} b^\dagger. \quad (6.18)$$

Their variances satisfy the general uncertainty relation

$$(\Delta x)^2 (\Delta p)^2 \geq \frac{1}{4} |\langle [x, p] \rangle|^2 = \frac{1}{4} |\langle [a, a^\dagger] \rangle|^2. \quad (6.19)$$

A state, for which there is equality in (6.19), is said to satisfy the minimum uncertainty property. If the vacuum state satisfies the minimum uncertainty property, then the deformed quadrature x (resp. p) is said to be squeezed in a given state if $X \equiv (\Delta x)^2 / (\Delta x)_0^2$ (resp. $P \equiv (\Delta p)^2 / (\Delta p)_0^2$) is less than one.

We shall now proceed to study both types of squeezing properties for the CS $|z; \mu; \alpha\rangle$ and $|z\rangle$.

6.2.1 Squeezing Properties of the Coherent States $|z; \mu; \alpha\rangle$

In any state belonging to $\mathcal{F}_\mu$, the variances of the deformed quadratures satisfy the uncertainty relation

$$(\Delta x)^2 (\Delta p)^2 \geq \frac{\lambda^2}{4} (\bar{\beta}_{\mu+1} - \bar{\beta}_\mu)^2, \quad (6.20)$$

where the right-hand side may be less than the conventional value $1/4$ [33]. They can be expressed as

$$(\Delta x)^2 = \langle H_0 \rangle + \delta_{\lambda,2} \langle J_+ + J_- \rangle, \quad (\Delta p)^2 = \langle H_0 \rangle - \delta_{\lambda,2} \langle J_+ + J_- \rangle. \quad (6.21)$$

Since in $\mathcal{F}_\mu$ the role of the vacuum state is played by the number state $|\mu\rangle = |0; \mu; \alpha\rangle$, and $\langle H_0 \rangle \geq \langle H_0 \rangle_0$, it is obvious that there cannot be any squeezing for $\lambda \geq 3$.¹

For $\lambda = 2$ and $\alpha = 0$, we obtain

$$(\Delta x)^2 = \langle N \rangle + \bar{\beta}_1 + \text{Re } z, \quad (\Delta p)^2 = \langle N \rangle + \bar{\beta}_1 - \text{Re } z, \quad (6.22)$$

$$(\Delta x)_0^2 = (\Delta p)_0^2 = \bar{\beta}_1 + \mu = \bar{\beta}_{\mu+1} + \bar{\beta}_\mu, \quad (6.23)$$

where $\langle N \rangle$ is given by

$$\begin{aligned}\langle N \rangle &= \frac{2y}{\beta_1} \Phi_0^1(y; 0; 0) \quad \text{if } \mu = 0, \\ &= 1 + 2\bar{\beta}_1 [\Phi_1^0(y; 1; 0) - 1] \quad \text{if } \mu = 1.\end{aligned}\tag{6.24}$$

Comparing (6.20) and (6.23), we see that $|\mu\rangle$ satisfies the minimum uncertainty property only for $\mu = 0$, to which we shall restrict ourselves.

From (6.22) and (6.23), it is obvious that X and P are related to each other by the transformation $\text{Re } z \rightarrow -\text{Re } z$ and that the maximum squeezing in x is achieved for real, negative values of z . So we only consider X for such values. As shown in Fig. 5a, a large squeezing effect over the whole range of real, negative values of z is obtained for $\alpha_0 > 0$ (i.e., $\bar{\beta}_1 > 1/2$).

For the conventional quadratures, the counterpart of (6.21) is

$$(\Delta x_b)^2 = \langle N \rangle + \frac{1}{2} + \delta_{\lambda,2} \text{Re } z \left\langle \left(\frac{(N+1)(N+2)}{F(N+1)F(N+2)} \right)^{1/2} \right\rangle, \tag{6.25}$$

$$(\Delta p_b)^2 = \langle N \rangle + \frac{1}{2} - \delta_{\lambda,2} \text{Re } z \left\langle \left(\frac{(N+1)(N+2)}{F(N+1)F(N+2)} \right)^{1/2} \right\rangle, \tag{6.26}$$

where use has been made of (6.18). There is no squeezing for $\lambda \geq 3$ again.

For $\lambda = 2$ and $\mu = \alpha = 0$, we obtain from (2.7), (2.10), (3.11), and (3.12),

$$\left\langle \left(\frac{(N+1)(N+2)}{F(N+1)F(N+2)} \right)^{1/2} \right\rangle = [{}_0F_1(\bar{\beta}_1; y)]^{-1} \sum_{k=0}^{\infty} \left(\frac{2k+1}{2k+2\bar{\beta}_1} \right)^{1/2} \frac{y^k}{k!(\bar{\beta}_1)_k}, \tag{6.27}$$

where $y = |z|^2/4$. The previous remarks about X and P remain valid. For real, negative values of z , the behaviour of X in terms of $-\text{Re } z$ is shown in Fig. 5b. Comparing with Fig. 5a, we observe more or less similar squeezing properties to those corresponding to dressed photons.

6.2.2 Squeezing Properties of the Coherent States $|z\rangle$

From (2.2) and (6.17), it follows that in any state belonging to $\mathcal{F}$, the variances of the deformed quadratures can be expressed as

$$(\Delta x)^2 = \langle H_0 \rangle - \langle a \rangle \langle a^\dagger \rangle + \frac{1}{2} [(\Delta a)^2 + (\Delta a^\dagger)^2], \tag{6.28}$$

$$(\Delta p)^2 = \langle H_0 \rangle - \langle a \rangle \langle a^\dagger \rangle - \frac{1}{2} [(\Delta a)^2 + (\Delta a^\dagger)^2]. \tag{6.29}$$

For the CS $|z\rangle$, $\langle a \rangle = z$, $\langle a^\dagger \rangle = z^*$, and $(\Delta a)^2 = (\Delta a^\dagger)^2 = 0$, so that $(\Delta x)^2 = (\Delta p)^2 = \langle H_0 \rangle - |z|^2$. Furthermore, since $[a, a^\dagger] = \{a, a^\dagger\} - 2a^\dagger a$, we also find $\frac{1}{2} \langle [a, a^\dagger] \rangle = \langle H_0 \rangle - |z|^2$. We therefore conclude that the CS $|z\rangle$ satisfy the minimum uncertainty property and that in such states

$$(\Delta x)^2 = (\Delta p)^2 = \frac{\lambda}{2} [\mathcal{N}(|z|)]^{-1} \sum_{\mu=0}^{\lambda-1} (\bar{\beta}_{\mu+1} - \bar{\beta}_\mu) N_\mu^{(0)}(|\omega|) \frac{t^\mu}{\prod_{\nu=1}^\mu \bar{\beta}_\nu}, \quad (6.30)$$

where $t = |z|^2/\lambda$, $|\omega| = |z|^\lambda$. For the vacuum state $|0\rangle$, Eq. (6.30) reduces to

$$(\Delta x)_0^2 = (\Delta p)_0^2 = \frac{\lambda}{2} \bar{\beta}_1, \quad (6.31)$$

which is less than the conventional value $1/2$ for $\bar{\beta}_1 < \lambda^{-1}$ (i.e., $\alpha_0 < 0$). The variances of the deformed quadratures x and p are simultaneously lower than their common vacuum value if $X = P < 1$.

For $\lambda = 2$ corresponding to the paraboson CS [35], $X = P$ can be rewritten in terms of Bessel modified functions I_ν as

$$X = P = 1 + \frac{1 - 2\bar{\beta}_1}{\bar{\beta}_1} R(t), \quad (6.32)$$

where $R(t)$ is defined in (6.14). They behave as $1 + (1 - 2\bar{\beta}_1)|z|^2/(2\bar{\beta}_1^2)$ or $(2\bar{\beta}_1)^{-1}$ according to whether $|z| \ll 1$ or $|z| \gg 1$. In these limits, we therefore get $X = P < 1$ if $\bar{\beta}_1 > 1/2$ (i.e., $\alpha_0 > 0$). In Fig. 6a, it is shown that this result is actually valid over the whole range of $|z|$ values.

For $\lambda \geq 3$, X and P behave as $1 + (\bar{\beta}_2 - 2\bar{\beta}_1)|z|^2/(\lambda\bar{\beta}_1^2)$ or $(\lambda\bar{\beta}_1)^{-1}$ according to whether $|z| \ll 1$ or $|z| \gg 1$. Values of $\bar{\beta}_1$ greater than both $1/\lambda$ and $\bar{\beta}_2/2$ (i.e., $\alpha_0 > 0$ and $\alpha_1 < \alpha_0$) therefore favour small values of $X = P$. This is confirmed numerically, as shown in Fig. 6b for $\lambda = 3$. It should be noted however that there is a tendency for the behaviour of $X = P$ to become more complicated than in the $\lambda = 2$ case.

For the conventional quadratures, the equality of the variances obtained in (6.30) does not remain true. We obtain instead

$$\begin{aligned} (\Delta x_b)^2 &= \langle N \rangle + \frac{1}{2} - |z|^2 \left\langle \left(\frac{(N+1)(N+2)}{F(N+1)F(N+2)} \right)^{1/2} \right\rangle \\ &\quad + 2(\text{Re } z)^2 \left[\left\langle \left(\frac{(N+1)(N+2)}{F(N+1)F(N+2)} \right)^{1/2} \right\rangle^2 - \left\langle \left(\frac{N+1}{F(N+1)} \right)^{1/2} \right\rangle^2 \right], \end{aligned} \quad (6.33)$$

and $(\Delta p_b)^2$ given by a similar formula with $\text{Re } z$ replaced by $\text{Im } z$. Here

$$\langle N \rangle = \lambda [\mathcal{N}(|z|)]^{-1} S_1(|z|), \quad (6.34)$$

$$\begin{aligned} \left\langle \left(\frac{N+1}{F(N+1)} \right)^{1/2} \right\rangle &= [\mathcal{N}(|z|)]^{-1} \sum_{\mu=0}^{\lambda-1} \frac{t^\mu}{\prod_{\nu=1}^{\mu} \bar{\beta}_\nu} \sum_{k=0}^{\infty} \left(\frac{k\lambda + \mu + 1}{k\lambda + \lambda \bar{\beta}_{\mu+1}} \right)^{1/2} \\ &\quad \times \frac{t^{k\lambda}}{k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_\nu + 1)_k \right) \left(\prod_{\nu=\mu+1}^{\lambda-1} (\bar{\beta}_\nu)_k \right)}, \end{aligned} \quad (6.35)$$

$$\begin{aligned} \left\langle \left(\frac{(N+1)(N+2)}{F(N+1)F(N+2)} \right)^{1/2} \right\rangle &= [\mathcal{N}(|z|)]^{-1} \sum_{\mu=0}^{\lambda-1} \frac{t^\mu}{\prod_{\nu=1}^{\mu} \bar{\beta}_\nu} \\ &\quad \times \sum_{k=0}^{\infty} \left(\frac{(k\lambda + \mu + 1)(k\lambda + \mu + 2)}{(k\lambda + \lambda \bar{\beta}_{\mu+1})(k\lambda + \lambda \bar{\beta}_{\mu+2})} \right)^{1/2} \\ &\quad \times \frac{t^{k\lambda}}{k! \left(\prod_{\nu=1}^{\mu} (\bar{\beta}_\nu + 1)_k \right) \left(\prod_{\nu=\mu+1}^{\lambda-1} (\bar{\beta}_\nu)_k \right)}, \end{aligned} \quad (6.36)$$

where $S_1(|z|)$ is given in (6.11) and $t = |z|^2/\lambda$. Since X and P are related to each other by the transformation $\text{Re } z \leftrightarrow \text{Im } z$, it is enough to study the former.

For small $|z|$ values, the coefficient of $2(\text{Re } z)^2$ on the right-hand side of (6.33) behaves as $\left(\sqrt{2\bar{\beta}_1} - 1 \right) / (2\bar{\beta}_1)$ if $\lambda = 2$, and as $\left(\sqrt{2\bar{\beta}_1} - \sqrt{\bar{\beta}_2} \right) / \left(\lambda \bar{\beta}_1 \sqrt{\bar{\beta}_2} \right)$ if $\lambda \geq 3$. Hence, for such values, the maximum squeezing in x_b is achieved when z is real (resp. imaginary) if $\bar{\beta}_1 < 1/2$ (resp. $\bar{\beta}_1 > 1/2$) in the former case and if $\bar{\beta}_1 < \bar{\beta}_2/2$ (resp. $\bar{\beta}_1 > \bar{\beta}_2/2$) in the latter.

A numerical study does confirm this trend. In Fig. 7, X is plotted against $\text{Re } z$ or $\text{Im } z$ for $\lambda = 2$ and some values of $\bar{\beta}_1$ smaller or greater than $1/2$ (i.e., $\alpha_0 < 0$ or $\alpha_0 > 0$), respectively. In Fig. 8, the same is done for $\lambda = 3$ and some values of $(\bar{\beta}_1, \bar{\beta}_2)$ such that $\bar{\beta}_1 < \bar{\beta}_2/2$ or $\bar{\beta}_1 > \bar{\beta}_2/2$ (i.e., $\alpha_0 < \alpha_1$ or $\alpha_0 > \alpha_1$). We conclude that a substantial squeezing effect can be obtained for well-chosen parameters.

7 CONCLUSION

In the present paper, we introduced two new types of CS associated with the C_λ -extended oscillator. The first ones, which actually form a family of CS whose members are labelled by some index α , include as special cases both the Barut-Girardello and the Perelomov $\text{su}(1,1)$

CS for $\lambda = 2$, as well as the annihilation-operator CS of the C_λ -extended oscillator SGA for higher λ values. The second ones, which are eigenstates of the C_λ -extended oscillator annihilation operator, extend to higher λ values the paraboson CS, to which they reduce for $\lambda = 2$.

We showed that all these states satisfy a unity resolution relation in the C_λ -extended oscillator Fock space (or in some subspace thereof). It should be stressed that it is this property that makes them qualify as generalized CS and distinguishes them from the plethora of putative CS found in the literature. To arrive at this result for some of the CS considered here, it has been essential to restrict the range of the C_λ -extended oscillator parameters in such a way that the limit $\alpha_\mu \rightarrow 0$, corresponding to the standard harmonic oscillator, is excluded. The C_λ -extended oscillator parameters therefore play a regularizing role, similar to that of the parameter α_0 of the Calogero-Vasiliev algebra for the Perelomov $\text{su}(1,1)$ CS. It is indeed worth reminding that for the latter, the unity resolution relation invalid for the standard harmonic oscillator (i.e., for $\alpha_0 = 0$ or $\bar{\beta}_1 = 1/2$) becomes true for sufficiently large values of α_0 (i.e., for $\alpha_0 > 1$ or $\bar{\beta}_1 > 1$).

As a by-product of our study, we obtained Bargmann representations of the C_λ -extended oscillator Fock space. This allowed us to solve, albeit in the complex plane, the yet open problem of finding realizations of the C_λ -extended oscillator algebra in terms of differential operators.

Furthermore, we investigated some characteristics of our CS relevant to quantum optics, such as their statistical and squeezing properties, for a wide range of parameters and from both viewpoints of real and dressed photons. The nonclassical features of these states for some parameter ranges were clearly demonstrated.

The remarkable properties of the new CS defined in this paper provide a useful tool for theoretical investigation of model systems. Although their generation by nonlinear systems remains to be studied, we believe that these states may play an important role in quantum optics.

APPENDIX A: PROOF OF EQS. (4.13) AND (4.14)

In this Appendix, we prove that Eq. (4.13) provides us with a positive solution of (4.7) for any $r > 0$ and any $0 \leq \alpha \leq \lambda - \mu - 1$, if condition (4.14) is fulfilled.

From (4.12), it follows that Eq. (4.13) is valid for $\alpha = 0$ without any restriction. We will now show that if it is valid for $\alpha - 1$ under a condition similar to (4.14), then the same is true for α . For such a purpose, let us factorize $g^*(s)$, defined in (4.7), as

$$g^*(s) = g_1^*(s)g_2^*(s), \quad g_1^*(s) = \frac{\prod_{\nu=1}^{r+\alpha-1} \Gamma(s + b_\nu)}{\prod_{\nu=1}^{\alpha-1} \Gamma(s + a_\nu)}, \quad g_2^*(s) = \frac{\Gamma(s + b_{r+\alpha})}{\Gamma(s + a_\alpha)}. \quad (\text{A.1})$$

From the induction hypothesis and tables of inverse Mellin transforms [47], we get

$$g_1(y) = G_{\alpha-1, r+\alpha-1}^{r+\alpha-1, 0} \left(y \left| \begin{array}{c} a_1, a_2, \dots, a_{\alpha-1} \\ b_1, b_2, \dots, b_{r+\alpha-1} \end{array} \right. \right) \quad (\text{A.2})$$

and

$$g_2(y) = \begin{cases} \frac{y^{b_{r+\alpha}} (1-y)^{a_\alpha - b_{r+\alpha} - 1}}{\Gamma(a_\alpha - b_{r+\alpha})} & \text{if } 0 < y < 1 \\ 0 & \text{if } 1 < y < \infty \end{cases}, \quad (\text{A.3})$$

respectively. Equation (A.2) is valid if the set $\{b_1, b_2, \dots, b_{r+\alpha-1}\}$ contains some subset of $\alpha - 1$ elements $\{b_{i_1}, b_{i_2}, \dots, b_{i_{\alpha-1}}\}$ such that

$$a_1 > b_{i_1}, \quad a_2 > b_{i_2}, \quad \dots, \quad a_{\alpha-1} > b_{i_{\alpha-1}}, \quad (\text{A.4})$$

while Eq. (A.3) requires

$$a_\alpha > b_{r+\alpha}. \quad (\text{A.5})$$

The second form of $g(y)$ in (4.11) leads to the expression

$$g(y) = \frac{y^{b_{r+\alpha}}}{\Gamma(a_\alpha - b_{r+\alpha})} \int_y^\infty t^{-a_\alpha} (t - y)^{a_\alpha - b_{r+\alpha} - 1} G_{\alpha-1, r+\alpha-1}^{r+\alpha-1, 0} \left(t \left| \begin{array}{c} a_1, a_2, \dots, a_{\alpha-1} \\ b_1, b_2, \dots, b_{r+\alpha-1} \end{array} \right. \right) dt, \quad (\text{A.6})$$

which is a positive function if conditions (A.4) and (A.5) are satisfied. The change of variable $u = t/y$ transforms (A.6) into

$$g(y) = [\Gamma(a_\alpha - b_{r+\alpha})]^{-1} \int_1^\infty u^{-a_\alpha} (u - 1)^{a_\alpha - b_{r+\alpha} - 1} G_{\alpha-1, r+\alpha-1}^{r+\alpha-1, 0} \left(yu \left| \begin{array}{c} a_1, a_2, \dots, a_{\alpha-1} \\ b_1, b_2, \dots, b_{r+\alpha-1} \end{array} \right. \right) du, \quad (\text{A.7})$$

which is nothing else than the right-hand side of (4.13) when use is made of Eq. (7.811.3) of [50].

To extend conditions (A.4) and (A.5) to the more general ones given in (4.14), it is enough to note that $g^*(s)$, and hence $g(y)$, are symmetric under any permutation of the a_ν 's or the b_ν 's. In other words, considering other factorizations than that assumed in (A.1) would lead to the general condition (4.14), which completes the proof.

APPENDIX B: PROOF OF EQS. (4.19), (4.20), AND (4.21)

In this Appendix, we prove that Eqs. (4.19) and (4.20) provide us with a positive solution of (4.4) for any even λ , $\alpha = \lambda/2 \geq 2$, and $\mu = 0$, if condition (4.21) is fulfilled. For such a purpose, it is enough to consider the inversion problem of (4.7) for $r = 0$, any $\alpha \geq 2$, and $\mu = 0$.

Let us start with the $\alpha = 2$ case and factorize $g^*(s)$ as

$$g^*(s) = g_1^*(s)g_2^*(s), \quad g_i^*(s) = \frac{\Gamma(s + b_i)}{\Gamma(s + a_i)} \quad i = 1, 2. \quad (\text{B.1})$$

The inverse Mellin transforms of the factors are given by [47]

$$g_i(y) = \begin{cases} \frac{y^{b_i(1-y)^{a_i-b_i-1}}}{\Gamma(a_i-b_i)} & \text{if } 0 < y < 1 \\ 0 & \text{if } 1 < y < \infty \end{cases}, \quad (\text{B.2})$$

provided

$$a_i > b_i \quad i = 1, 2. \quad (\text{B.3})$$

The second form of $g(y)$ in (4.11) leads to the expression

$$g(y) = \frac{y^{b_2}}{\Gamma(a_1 - b_1)\Gamma(a_2 - b_2)} \int_y^1 t^{b_1-a_2}(1-t)^{a_1-b_1-1}(t-y)^{a_2-b_2-1} dt, \quad (\text{B.4})$$

if $0 < y < 1$ and to zero if $1 < y < \infty$. The right-hand side of (B.4) is clearly a positive function on $(0, 1)$ if condition (B.3) is fulfilled. Substituting $u = (1-t)/(1-y)$ into (B.4) transforms it into

$$g(y) = \frac{y^{b_2}(1-y)^{a_1+a_2-b_1-b_2-1}}{\Gamma(a_1-b_1)\Gamma(a_2-b_2)} \int_0^1 u^{a_1-b_1-1}(1-u)^{a_2-b_2-1}[1-(1-y)u]^{b_1-a_2} du, \quad (\text{B.5})$$

which is nothing else than

$$g(y) = \frac{y^{b_2}(1-y)^{a_1+a_2-b_1-b_2-1}}{\Gamma(a_1+a_2-b_1-b_2)} {}_2F_1(a_1-b_1, a_2-b_2; a_1+a_2-b_1-b_2; 1-y), \quad (\text{B.6})$$

when use is made of Eq. (3.197.3) of [50].

Had we considered the other possible factorization of $g^*(s)$, obtained from (B.1) by permuting b_1 and b_2 while leaving a_1, a_2 unchanged, we should have obtained

$$g(y) = \frac{y^{b_1}(1-y)^{a_1+a_2-b_1-b_2-1}}{\Gamma(a_1+a_2-b_1-b_2)} {}_2F_1(a_1-b_2, a_2-b_2; a_1+a_2-b_1-b_2; 1-y), \quad (\text{B.7})$$

if

$$a_1 > b_2, \quad a_2 > b_1. \quad (\text{B.8})$$

However the right-hand side of (B.7) can be rewritten in the form (B.6) by using Eq. (9.131.1) of [50]. We therefore conclude that Eq. (B.6) applies in both cases (B.3) and (B.8), thereby leading to (4.18).

For $\alpha > 2$, the generalization of (B.6) reads

$$\begin{aligned} g(y) &= \frac{y^{b_\alpha}}{\Gamma(a_1 - b_1) \left[\prod_{p=1}^{\alpha-2} \Gamma(a_{p+1} - b_p) \right]} \\ &\times \sum_{n_1 n_2 \dots n_{\alpha-2}} \frac{\prod_{p=1}^{\alpha-2} [\Gamma(a_{p+1} - b_p + n_p) \Gamma(\xi_{\alpha,p}(\mathbf{n}))]}{\left[\prod_{p=1}^{\alpha-3} \Gamma(\eta_{\alpha,p}(\mathbf{n})) \right] \Gamma(\zeta_\alpha(\mathbf{n})) \left(\prod_{p=1}^{\alpha-2} n_p! \right)} \\ &\times (1-y)^{\zeta_\alpha(\mathbf{n})-1} {}_2F_1(a_\alpha - b_{\alpha-1}, \eta_{\alpha,\alpha-2}(\mathbf{n}); \zeta_\alpha(\mathbf{n}); 1-y), \end{aligned} \quad (\text{B.9})$$

where

$$\begin{aligned} \xi_{\alpha,p}(\mathbf{n}) &= \sum_{q=1}^p (a_q - b_q + n_q), \\ \eta_{\alpha,p}(\mathbf{n}) &= \sum_{q=1}^{p+1} (a_q - b_q) + \sum_{q=1}^p n_q, \\ \zeta_\alpha(\mathbf{n}) &= \sum_{p=1}^{\alpha} (a_p - b_p) + \sum_{p=1}^{\alpha-2} n_p. \end{aligned} \quad (\text{B.10})$$

The positivity condition on $g(y)$ is now that there exists some permutation $(i_1, i_2, \dots, i_\alpha)$ of $(1, 2, \dots, \alpha)$ such that

$$a_1 > b_{i_1}, \quad a_2 > b_{i_2}, \quad \dots, \quad a_\alpha > b_{i_\alpha}. \quad (\text{B.11})$$

The proof of (B.9), (B.10), and (B.11) is by induction over α , using the factorization (A.1) of $g^*(s)$ for $r = 0$ and the convolution property (4.11). The intermediate steps are the same as in the $\alpha = 2$ case, once the hypergeometric function ${}_2F_1(a_{\alpha-1} - b_{\alpha-2}, \eta_{\alpha-1,\alpha-3}(\mathbf{n}); \zeta_{\alpha-1}(\mathbf{n}); (1-y)u)$ has been expanded into powers of $(1-y)u$. Finally, Eqs. (4.19), (4.20), and (4.21) are directly obtained from (B.9), (B.10), and (B.11) after permuting the b_ν 's in a convenient way and substituting (4.9) and (4.10).

APPENDIX C: HERMITICITY PROPERTIES OF SOME OPERATOR REALIZATIONS IN BARGMANN SPACE

In this Appendix, we consider the Hermiticity properties of some operators or some operator-valued matrices with respect to the scalar product (5.4) of $\mathcal{B}_\mu^{(\alpha)}$ or (5.18) of $\mathcal{B}^{(0)}$, respectively.

Let us start with the operators $\mathcal{J}_{+\mu}^{(\alpha)}$, $\mathcal{J}_{-\mu}^{(\alpha)}$, and $\mathcal{J}_{0\mu}^{(\alpha)}$, defined in (5.9)–(5.11), and prove that they satisfy relations similar to (2.14) with respect to (5.4). Considering the first relation in (2.14), we have to show that for any two functions $\psi_\mu^{(\alpha)}(z)$, $\psi_\mu^{\prime(\alpha)}(z) \in \mathcal{B}_\mu^{(\alpha)}$,

$$\int d^2z h_\mu^{(\alpha)}(y) [\mathcal{J}_{+\mu}^{(\alpha)} \psi_\mu^{(\alpha)}(z)]^* \psi_\mu^{(\alpha)}(z) = \int d^2z h_\mu^{(\alpha)}(y) [\psi_\mu^{\prime(\alpha)}(z)]^* \mathcal{J}_{-\mu}^{(\alpha)} \psi_\mu^{(\alpha)}(z). \quad (\text{C.1})$$

By substituting (5.10) into the right-hand side of (C.1), integrating by parts, and re-ordering some operators, we obtain

$$\begin{aligned} & \int d^2z h_\mu^{(\alpha)}(y) [\psi_\mu^{\prime(\alpha)}(z)]^* \mathcal{J}_{-\mu}^{(\alpha)} \psi_\mu^{(\alpha)}(z) \\ &= -\lambda^{\lambda-\alpha-1} \int d^2z \left\{ \frac{\partial}{\partial z} \left[\prod_{\nu=1}^{\mu} \left(-z \frac{\partial}{\partial z} + \bar{\beta}_\nu \right) \right] \left[\prod_{\nu=\mu+\alpha+1}^{\lambda-1} \left(-z \frac{\partial}{\partial z} + \bar{\beta}_\nu - 1 \right) \right] h_\mu^{(\alpha)}(y) \right\} \\ & \quad \times [\psi_\mu^{\prime(\alpha)}(z)]^* \psi_\mu^{(\alpha)}(z). \end{aligned} \quad (\text{C.2})$$

Since

$$\frac{\partial}{\partial z} h_\mu^{(\alpha)}(y) = \lambda^{2\alpha-\lambda} z \frac{d}{dy} h_\mu^{(\alpha)}(y), \quad z \frac{\partial}{\partial z} h_\mu^{(\alpha)}(y) = y \frac{d}{dy} h_\mu^{(\alpha)}(y), \quad (\text{C.3})$$

Eq. (C.2) becomes

$$\begin{aligned} & \int d^2z h_\mu^{(\alpha)}(y) [\psi_\mu^{\prime(\alpha)}(z)]^* \mathcal{J}_{-\mu}^{(\alpha)} \psi_\mu^{(\alpha)}(z) \\ &= (-1)^{\lambda-\alpha} \lambda^{\alpha-1} \int d^2z \left\{ z \frac{d}{dy} \left[\prod_{\nu=1}^{\mu} \left(y \frac{d}{dy} - \bar{\beta}_\nu \right) \right] \left[\prod_{\nu=\mu+\alpha+1}^{\lambda-1} \left(y \frac{d}{dy} - \bar{\beta}_\nu + 1 \right) \right] h_\mu^{(\alpha)}(y) \right\} \\ & \quad \times [\psi_\mu^{\prime(\alpha)}(z)]^* \psi_\mu^{(\alpha)}(z). \end{aligned} \quad (\text{C.4})$$

By proceeding similarly, the left-hand side of (C.1) is transformed into

$$\begin{aligned} & \int d^2z h_\mu^{(\alpha)}(y) [\mathcal{J}_{+\mu}^{(\alpha)} \psi_\mu^{(\alpha)}(z)]^* \psi_\mu^{(\alpha)}(z) \\ &= (-1)^\alpha \lambda^{\alpha-1} \int d^2z \left\{ z \left[\prod_{\nu=\mu+1}^{\mu+\alpha} \left(y \frac{d}{dy} - \bar{\beta}_\nu + 2 \right) \right] h_\mu^{(\alpha)}(y) \right\} [\psi_\mu^{\prime(\alpha)}(z)]^* \psi_\mu^{(\alpha)}(z). \end{aligned} \quad (\text{C.5})$$

Equation (C.1) is therefore equivalent to the following differential equation for $h_\mu^{(\alpha)}(y)$,

$$\left\{ (-1)^\alpha \left[\prod_{\nu=\mu+1}^{\mu+\alpha} \left(y \frac{d}{dy} - \bar{\beta}_\nu + 2 \right) \right] - (-1)^{\lambda-\alpha} \frac{d}{dy} \left[\prod_{\nu=1}^{\mu} \left(y \frac{d}{dy} - \bar{\beta}_\nu \right) \right] \right. \\ \left. \times \left[\prod_{\nu=\mu+\alpha+1}^{\lambda-1} \left(y \frac{d}{dy} - \bar{\beta}_\nu + 1 \right) \right] \right\} h_\mu^{(\alpha)}(y) = 0. \quad (\text{C.6})$$

After multiplication by y , Eq. (C.6) is nothing else than the differential equation satisfied by the Meijer G -function given in (4.15) [45]. This completes the proof of (C.1).

The proof of the second relation in (2.14) is easily carried out by using only integrations by parts and relations such as (C.3).

Let us now turn ourselves to the operator-valued matrices $\mathcal{A}^{\dagger(0)}$ and $\mathcal{A}^{(0)}$, defined in (5.23) and (5.24), respectively, and show that they are Hermitian conjugate of one another with respect to (5.18), i.e.,

$$\sum_{\mu, \mu'=0}^{\lambda-1} \int d^2 z h_\mu^{(0)}(y) \left[\mathcal{A}_{\mu\mu'}^{\dagger(0)} \psi_{\mu'}^{(0)}(z) \right]^* \psi_\mu^{(0)}(z) = \sum_{\mu, \mu'=0}^{\lambda-1} \int d^2 z h_\mu^{(0)}(y) \left[\psi_\mu^{(0)}(z) \right]^* \mathcal{A}_{\mu\mu'}^{(0)} \psi_{\mu'}^{(0)}(z). \quad (\text{C.7})$$

Substituting the explicit expressions of $\mathcal{A}_{\mu\mu'}^{\dagger(0)}$ into the left-hand side of (C.7) directly leads to

$$\sum_{\mu, \mu'=0}^{\lambda-1} \int d^2 z h_\mu^{(0)}(y) \left[\mathcal{A}_{\mu\mu'}^{\dagger(0)} \psi_{\mu'}^{(0)}(z) \right]^* \psi_\mu^{(0)}(z) \\ = \left[\lambda^{\lambda-1} \left(\prod_{\nu=1}^{\lambda-1} \bar{\beta}_\nu \right) \right]^{-1/2} \int d^2 z h_0^{(0)}(y) \left[z \psi_{\lambda-1}'^{(0)}(z) \right]^* \psi_0^{(0)}(z) \\ + \sum_{\mu=0}^{\lambda-2} \sqrt{\lambda \bar{\beta}_{\mu+1}} \int d^2 z h_{\mu+1}^{(0)}(y) \left[\psi_\mu^{(0)}(z) \right]^* \psi_{\mu+1}^{(0)}(z). \quad (\text{C.8})$$

After substituting the explicit expressions of $\mathcal{A}_{\mu\mu'}^{(0)}$ into the right-hand side of (C.7), integrating by parts, re-ordering some operators, and using (C.3) for $\alpha = 0$, we get

$$\sum_{\mu, \mu'=0}^{\lambda-1} \int d^2 z h_\mu^{(0)}(y) \left[\psi_\mu^{(0)}(z) \right]^* \mathcal{A}_{\mu\mu'}^{(0)} \psi_{\mu'}^{(0)}(z) \\ = - \left[\lambda^{1-\lambda} \left(\prod_{\nu=1}^{\lambda-1} \bar{\beta}_\nu \right) \right]^{1/2} \int d^2 z \left[\frac{d}{dy} h_{\lambda-1}^{(0)}(y) \right] \left[z \psi_{\lambda-1}'^{(0)}(z) \right]^* \psi_0^{(0)}(z) \\ - \sum_{\mu=0}^{\lambda-2} \sqrt{\frac{\lambda}{\bar{\beta}_{\mu+1}}} \int d^2 z \left[\left(y \frac{d}{dy} - \bar{\beta}_{\mu+1} + 1 \right) h_\mu^{(0)}(y) \right] \left[\psi_\mu^{(0)}(z) \right]^* \psi_{\mu+1}^{(0)}(z). \quad (\text{C.9})$$

Equation (C.7) is therefore equivalent to the following system of λ differential equations for $h_\mu^{(0)}(y)$, $\mu = 0, 1, \dots, \lambda - 1$,

$$\left(y \frac{d}{dy} - \bar{\beta}_{\mu+1} + 1\right) h_\mu^{(0)}(y) = -\bar{\beta}_{\mu+1} h_{\mu+1}^{(0)}(y), \quad \mu = 0, 1, \dots, \lambda - 2, \quad (\text{C.10})$$

$$\frac{d}{dy} h_{\lambda-1}^{(0)}(y) = -\left(\prod_{\nu=1}^{\lambda-1} \bar{\beta}_\nu\right)^{-1} h_0^{(0)}(y). \quad (\text{C.11})$$

By taking (4.5) and (4.15) into account, this leads to the system

$$\begin{aligned} & \left(y \frac{d}{dy} - \bar{\beta}_{\mu+1} + 1\right) G_{0\lambda}^{\lambda 0}(y|0, \bar{\beta}_1, \dots, \bar{\beta}_\mu, \bar{\beta}_{\mu+1} - 1, \dots, \bar{\beta}_{\lambda-1} - 1) \\ &= -G_{0\lambda}^{\lambda 0}(y|0, \bar{\beta}_1, \dots, \bar{\beta}_{\mu+1}, \bar{\beta}_{\mu+2} - 1, \dots, \bar{\beta}_{\lambda-1} - 1), \quad \mu = 0, 1, \dots, \lambda - 2, \end{aligned} \quad (\text{C.12})$$

$$\frac{d}{dy} G_{0\lambda}^{\lambda 0}(y|0, \bar{\beta}_1, \dots, \bar{\beta}_{\lambda-1}) = -G_{0\lambda}^{\lambda 0}(y|0, \bar{\beta}_1 - 1, \dots, \bar{\beta}_{\lambda-1} - 1), \quad (\text{C.13})$$

to be satisfied by Meijer G -functions.

To prove (C.12), we successively use Eq. (5.3.1.9) of [45], the symmetry of the G -function in its parameters, Eq. (5.3.1.13) of the same, and the symmetry again to rewrite its left-hand side as

$$\begin{aligned} & \left(y \frac{d}{dy} - \bar{\beta}_{\mu+1} + 1\right) G_{0\lambda}^{\lambda 0}(y|0, \bar{\beta}_1, \dots, \bar{\beta}_\mu, \bar{\beta}_{\mu+1} - 1, \dots, \bar{\beta}_{\lambda-1} - 1) \\ &= -\left(u \frac{d}{du} + \bar{\beta}_{\mu+1} - 1\right) G_{\lambda 0}^{0\lambda}(u|1, 1 - \bar{\beta}_1, \dots, 1 - \bar{\beta}_\mu, 2 - \bar{\beta}_{\mu+1}, \dots, 2 - \bar{\beta}_{\lambda-1}) \\ &= -\left(u \frac{d}{du} + \bar{\beta}_{\mu+1} - 1\right) G_{\lambda 0}^{0\lambda}(u|2 - \bar{\beta}_{\mu+1}, 1, 1 - \bar{\beta}_1, \dots, 1 - \bar{\beta}_\mu, 2 - \bar{\beta}_{\mu+2}, \dots, 2 - \bar{\beta}_{\lambda-1}) \\ &= -G_{\lambda 0}^{0\lambda}(u|1 - \bar{\beta}_{\mu+1}, 1, 1 - \bar{\beta}_1, \dots, 1 - \bar{\beta}_\mu, 2 - \bar{\beta}_{\mu+2}, \dots, 2 - \bar{\beta}_{\lambda-1}) \\ &= -G_{\lambda 0}^{0\lambda}(u|1, 1 - \bar{\beta}_1, \dots, 1 - \bar{\beta}_{\mu+1}, 2 - \bar{\beta}_{\mu+2}, \dots, 2 - \bar{\beta}_{\lambda-1}), \end{aligned} \quad (\text{C.14})$$

where $u = 1/y$. Applying Eq. (5.3.1.9) of [45] then leads to the right-hand side of (C.12).

Similarly, the left-hand side of (C.13) can be transformed into

$$\begin{aligned} & \frac{d}{dy} G_{0\lambda}^{\lambda 0}(y|0, \bar{\beta}_1, \dots, \bar{\beta}_{\lambda-1}) = -\frac{1}{y} u \frac{d}{du} G_{\lambda 0}^{0\lambda}(u|1, 1 - \bar{\beta}_1, \dots, 1 - \bar{\beta}_{\lambda-1}) \\ &= -\frac{1}{y} G_{\lambda 0}^{0\lambda}(u|0, 1 - \bar{\beta}_1, \dots, 1 - \bar{\beta}_{\lambda-1}) = -\frac{1}{y} G_{0\lambda}^{\lambda 0}(y|1, \bar{\beta}_1, \dots, \bar{\beta}_{\lambda-1}), \end{aligned} \quad (\text{C.15})$$

which gives the right-hand side of (C.13) when using Eq. (5.3.1.8) of [45].

ACKNOWLEDGMENTS

The author is a Research Director of the National Fund for Scientific Research (FNRS), Belgium.

References

- [1] R. J. Glauber, *Phys. Rev.* **130** (1963), 2529; *Phys. Rev.* **131** (1963), 2766.
- [2] J. R. Klauder, *J. Math. Phys.* **4** (1963), 1055; *J. Math. Phys.* **4** (1963), 1058.
- [3] E. C. G. Sudarshan, *Phys. Rev. Lett.* **10** (1963), 277.
- [4] E. Schrödinger, *Naturwiss.* **14** (1926), 664.
- [5] A. O. Barut and L. Girardello, *Commun. Math. Phys.* **21** (1971), 41.
- [6] A. M. Perelomov, *Commun. Math. Phys.* **26** (1972), 222; R. Gilmore, *Ann. Phys.* (N.Y.) **74** (1972), 391.
- [7] C. Aragone, G. Guerri, S. Salamó, and J. L. Tani, *J. Phys. A* **7** (1974), L149; M. M. Nieto and L. M. Simmons, Jr., *Phys. Rev. Lett.* **41** (1978), 207.
- [8] J. R. Klauder and B. S. Skagerstam, “Coherent States — Applications in Physics and Mathematics,” World Scientific, Singapore, 1985.
- [9] A. M. Perelomov, “Generalized Coherent States and Their Applications,” Springer, Berlin, 1986.
- [10] W.-M. Zhang, D. H. Feng, and R. Gilmore, *Rev. Mod. Phys.* **62** (1990), 867.
- [11] H. J. Kimble, M. Dagenais, and L. Mandel, *Phys. Rev. Lett.* **39** (1977), 691.
- [12] R. Short and L. Mandel, *Phys. Rev. Lett.* **51** (1983), 384; M. C. Teich and B. E. A. Saleh, *J. Opt. Soc. Am. B* **2** (1985), 275.
- [13] C. K. Hong and L. Mandel, *Phys. Rev. Lett.* **54** (1985), 323; R. E. Slusher, L. W. Hollberg, B. Yurke, J. C. Mertz, and J. F. Valley, *Phys. Rev. Lett.* **55** (1985), 2409; L.-A. Wu, H. J. Kimble, J. L. Hall, and H. Wu, *Phys. Rev. Lett.* **57** (1986), 2520.
- [14] V. V. Dodonov, I. A. Malkin, and V. I. Man’ko, *Physica* **72** (1974), 597.
- [15] V. Bužek, I. Jex, and Tran Quang, *J. Mod. Opt.* **37** (1990), 159.

- [16] G. M. D'Ariano, M. G. Rasetti, J. Katriel, and A. I. Solomon, in "Squeezed and Nonclassical Light" (P. Tombesi and E. R. Pike, Eds.), p. 301, Plenum, New York, 1989; P. Shanta, S. Chaturvedi, V. Srinivasan, G. S. Agarwal, and C. L. Mehta, *Phys. Rev. Lett.* **72** (1994), 1447.
- [17] C. Daskaloyannis, *J. Phys. A* **24** (1991), L789.
- [18] A. I. Solomon, *Phys. Lett. A* **196** (1994), 29; J. Katriel and A. I. Solomon, *Phys. Rev. A* **49** (1994), 5149; A. I. Solomon, in "Fifth Int. Conf. on Squeezed States and Uncertainty Relations, Balatonfured, Hungary, May 27–31, 1997" (D. Han, J. Janszky, Y. S. Kim, and V. I. Man'ko, Eds.), p. 157, NASA Goddard Space Flight Center, Greenbelt, 1998.
- [19] J. Katriel and C. Quesne, *J. Math. Phys.* **37** (1996), 1650.
- [20] V. I. Man'ko, G. Marmo, F. Zaccaria, and E. C. G. Sudarshan, *Phys. Scr.* **55** (1997), 528.
- [21] P. Shanta, S. Chaturvedi, V. Srinivasan, and R. Jagannathan, *J. Phys. A* **27** (1994), 6433.
- [22] R. L. de Matos Filho and W. Vogel, *Phys. Rev. A* **54** (1996), 4560.
- [23] D. J. Fernández and V. Hussin, *J. Phys. A* **32** (1999), 3603.
- [24] S. Mancini, *Phys. Lett. A* **233** (1997), 291; S. Sivakumar, *Phys. Lett. A* **250** (1998), 257.
- [25] X.-M. Liu, *J. Phys. A* **32** (1999), 8685.
- [26] G. Junker and P. Roy, *Phys. Lett. A* **257** (1999), 113.
- [27] C. Quesne and N. Vansteenkiste, *Phys. Lett. A* **240** (1998), 21.
- [28] M. Daoud and M. Kibler, A fractional supersymmetric oscillator and its coherent states, preprint, math-ph/9912024.
- [29] C. Quesne and N. Vansteenkiste, *Helv. Phys. Acta* **72** (1999), 71.
- [30] C. Quesne and N. Vansteenkiste, *Int. J. Theor. Phys.* **39** (2000), 1175.

- [31] M. A. Vasiliev, *Int. J. Mod. Phys. A* **6** (1991), 1115.
- [32] T. Brzeziński, I. L. Egusquiza, and A. J. Macfarlane, *Phys. Lett. B* **311** (1993), 202.
- [33] C. Quesne, *Phys. Lett. A* **272** (2000), 313; Erratum, *Phys. Lett. A* **275** (2000), 313.
- [34] J.-M. Sixdeniers, K. A. Penson, and A. I. Solomon, *J. Phys. A* **32** (1999), 7543.
- [35] J. K. Sharma, C. L. Mehta, and E. C. G. Sudarshan, *J. Math. Phys.* **19** (1978), 2089;
J. K. Sharma, C. L. Mehta, N. Mukunda, and E. C. G. Sudarshan, *J. Math. Phys.* **22**
(1981), 78.
- [36] S. Chaturvedi and V. Srinivasan, *Phys. Rev. A* **44** (1991), 8024; A. J. Macfarlane, *J. Math. Phys.* **35** (1994), 1054.
- [37] Y. Ohnuki and S. Kamefuchi, “Quantum Field Theory and Parastatistics,” Springer, Berlin, 1982.
- [38] V. Bargmann, *Commun. Pure Appl. Math.* **14** (1961), 187.
- [39] J. Deenen and C. Quesne, *J. Math. Phys.* **25** (1984), 2354; D. J. Rowe, *J. Math. Phys.* **25** (1984), 2662; K. T. Hecht, “The Vector Coherent State Method and Its Application to Problems of Higher Symmetries,” Springer, Berlin, 1987.
- [40] A. P. Polychronakos, *Mod. Phys. Lett. A* **5** (1990), 2325; M. Roček, *Phys. Lett. B* **255** (1991), 554.
- [41] N. I. Akhiezer, “The Classical Moment Problem and Some Related Questions in Analysis,” Oliver and Boyd, London, 1965.
- [42] I. N. Sneddon, “The Use of Integral Transforms,” McGraw-Hill, New York, 1974.
- [43] K. A. Penson and A. I. Solomon, *J. Math. Phys.* **40** (1999), 2354.
- [44] J.-M. Sixdeniers and K. A. Penson, *J. Phys. A* **33** (2000), 2907.
- [45] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, “Higher Transcendental Functions,” Vol. 1, McGraw-Hill, New York, 1953.

- [46] P. Appell and J. Kampé de Fériet, “Fonctions Hypergéométriques et Hypersphériques. Polynômes d’Hermite,” Gauthier-Villars, Paris, 1926.
- [47] A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev, “Integrals and Series,” Vol. 3, Gordon and Breach, New York, 1990.
- [48] J. Peřina, “Quantum Statistics of Linear and Nonlinear Optical Phenomena,” Reidel, Dordrecht, 1984.
- [49] L. Mandel and E. Wolf, “Optical Coherence and Quantum Optics,” Cambridge University Press, Cambridge, 1995.
- [50] I. S. Gradshteyn and I. M. Ryzhik, “Table of Integrals, Series, and Products,” Academic Press, New York, 1980.

FOOTNOTES

¹ This is only valid for the second-order squeezing. In [33], we actually showed that there is a small fourth-order squeezing effect for $\lambda = 4$, $\mu = 0$, $\alpha = 0$.

FIGURE CAPTIONS

FIG. 1. The weight function $h_0^{(1)}(y)$ in terms of y for $\lambda = 3$. The parameter values are $\bar{\beta}_1 = 4/3$, $\bar{\beta}_2 = 2/3$, i.e., $\alpha_0 = -\alpha_1 = 3$ (solid line), or $\bar{\beta}_1 = \bar{\beta}_2 = 4/3$, i.e., $\alpha_0 = 3$, $\alpha_1 = -1$ (dashed line).

FIG. 2. The weight function $h_0^{(2)}(y)$ in terms of y for $\lambda = 4$ and $\bar{\beta}_1 = 3/2$, i.e., $\alpha_0 = 5$. The other parameter values are: (a) $\bar{\beta}_2 = 3/2$, $\bar{\beta}_3 = 5/4$, i.e., $\alpha_1 = -1$, $\alpha_2 = -2$ (solid line), $\bar{\beta}_2 = 7/4$, $\bar{\beta}_3 = 5/4$, i.e., $\alpha_1 = 0$, $\alpha_2 = -3$ (dashed line), or $\bar{\beta}_2 = 2$, $\bar{\beta}_3 = 5/4$, i.e., $\alpha_1 = 1$, $\alpha_2 = -4$ (dot-dashed line); (b) $\bar{\beta}_2 = 1$, $\bar{\beta}_3 = 3/4$, i.e., $\alpha_1 = -3$, $\alpha_2 = -2$ (solid line), $\bar{\beta}_2 = 5/4$, $\bar{\beta}_3 = 3/4$, i.e., $\alpha_1 = -2$, $\alpha_2 = -3$ (dashed line), or $\bar{\beta}_2 = 3/2$, $\bar{\beta}_3 = 3/4$, i.e., $\alpha_1 = -1$, $\alpha_2 = -4$ (dot-dashed line).

FIG. 3. Mandel's parameter Q as a function of $|z| \equiv r$ for the CS $|z; 1; \mu\rangle$, $\lambda = 3$, and various parameters: (a) $\mu = 0$ and $\bar{\beta}_1 = 1/3$, $\bar{\beta}_2 = 2/3$, i.e., $\alpha_0 = \alpha_1 = 0$ (solid line), $\bar{\beta}_1 = 1$, $\bar{\beta}_2 = 1/10$, i.e., $\alpha_0 = 2$, $\alpha_1 = -37/10$ (dashed line), $\bar{\beta}_1 = 1$, $\bar{\beta}_2 = 1/100$, i.e., $\alpha_0 = 2$, $\alpha_1 = -397/100$ (dotted line), or $\bar{\beta}_1 = 1/10$, $\bar{\beta}_2 = 2/3$, i.e., $\alpha_0 = -\alpha_1 = -7/10$ (dot-dashed line); (b) $\mu = 1$, $\bar{\beta}_1 = 1/3$, i.e., $\alpha_0 = 0$, and $\bar{\beta}_2 = 2/3$, i.e., $\alpha_1 = 0$ (solid line), $\bar{\beta}_2 = 1/10$, i.e., $\alpha_1 = -17/10$ (dashed line), $\bar{\beta}_2 = 1/50$, i.e., $\alpha_1 = -97/50$ (dotted line), or $\bar{\beta}_2 = 2$, i.e., $\alpha_1 = 4$ (dot-dashed line).

FIG. 4. Mandel's parameter Q as a function of $|z| \equiv r$ for the CS $|z\rangle$ and various parameters: (a) $\lambda = 2$ and $\bar{\beta}_1 = 1/90$, i.e., $\alpha_0 = -44/45$ (solid line), $\bar{\beta}_1 = 1/4$, i.e., $\alpha_0 = -1/2$ (dashed line), $\bar{\beta}_1 = 1$, i.e., $\alpha_0 = 1$ (dotted line), or $\bar{\beta}_1 = 10$, i.e., $\alpha_0 = 19$ (dot-dashed line); (b) $\lambda = 3$ and $\bar{\beta}_1 = 1/100$, $\bar{\beta}_2 = 2/3$, i.e., $\alpha_0 = -\alpha_1 = -97/100$ (solid line), $\bar{\beta}_1 = 1/3$, $\bar{\beta}_2 = 10$, i.e., $\alpha_0 = 0$, $\alpha_1 = 28$ (dashed line), $\bar{\beta}_1 = 2/3$, $\bar{\beta}_2 = 1/100$, i.e., $\alpha_0 = 1$, $\alpha_1 = -297/100$ (dotted line), or $\bar{\beta}_1 = 10$, $\bar{\beta}_2 = 1/3$, i.e., $\alpha_0 = 29$, $\alpha_1 = -30$ (dot-dashed line).

FIG. 5. X as a function of $-\text{Re } z$ for the CS $|z; 0; 0\rangle$, $\lambda = 2$, $\text{Im } z = 0$, and (a) the case of dressed photons, or (b) the case of real photons. The parameter values are $\bar{\beta}_1 = 1/2$, i.e., $\alpha_0 = 0$ (solid lines), $\bar{\beta}_1 = 3/10$, i.e., $\alpha_0 = -2/5$ (dashed lines), $\bar{\beta}_1 = 1$, i.e., $\alpha_0 = 1$ (dotted lines), or $\bar{\beta}_1 = 2$, i.e., $\alpha_0 = 3$ (dot-dashed lines).

FIG. 6. The ratio $X = (\Delta x)^2/(\Delta x)_0^2$ as a function of $|z| \equiv r$ for the CS $|z\rangle$ and various parameters: (a) $\lambda = 2$ and $\bar{\beta}_1 = 10$, i.e., $\alpha_0 = 19$ (solid line), $\bar{\beta}_1 = 2$, i.e., $\alpha_0 = 3$ (dashed line), $\bar{\beta}_1 = 3/4$, i.e., $\alpha_0 = 1/2$ (dotted line), or $\bar{\beta}_1 = 3/10$, i.e., $\alpha_0 = -2/5$ (dot-dashed line); (b) $\lambda = 3$ and $\bar{\beta}_1 = 2$, $\bar{\beta}_2 = 1/20$, i.e., $\alpha_0 = 5$, $\alpha_1 = -137/20$ (solid line), $\bar{\beta}_1 = \bar{\beta}_2 = 2/3$, i.e., $\alpha_0 = -\alpha_1 = 1$ (dashed line), $\bar{\beta}_1 = 2$, $\bar{\beta}_2 = 5$, i.e., $\alpha_0 = 5$, $\alpha_1 = 8$ (dotted line), or $\bar{\beta}_1 = 1/4$, $\bar{\beta}_2 = 1/8$, i.e., $\alpha_0 = -1/4$, $\alpha_1 = -11/8$ (dot-dashed line).

FIG. 7. The ratio $X = (\Delta x_b)^2/(\Delta x_b)_0^2$ for $\lambda = 2$ as a function of (a) $\text{Re } z$ for $\text{Im } z = 0$ and $\bar{\beta}_1 = 1/4$, i.e., $\alpha_0 = -1/2$ (solid line), $\bar{\beta}_1 = 1/10$, i.e., $\alpha_0 = -4/5$ (dashed line), $\bar{\beta}_1 = 1/40$, i.e., $\alpha_0 = -19/20$ (dotted line), or $\bar{\beta}_1 = 1/100$, i.e., $\alpha_0 = -49/50$ (dot-dashed line); (b) $\text{Im } z$ for $\text{Re } z = 0$ and $\bar{\beta}_1 = 1$, i.e., $\alpha_0 = 1$ (solid line), $\bar{\beta}_1 = 4$, i.e., $\alpha_0 = 7$ (dashed line), $\bar{\beta}_1 = 10$, i.e., $\alpha_0 = 19$ (dotted line), or $\bar{\beta}_1 = 40$, i.e., $\alpha_0 = 79$ (dot-dashed line).

FIG. 8. The ratio $X = (\Delta x_b)^2/(\Delta x_b)_0^2$ for $\lambda = 3$ as a function of (a) $\text{Re } z$ for $\text{Im } z = 0$ and $\bar{\beta}_1 = 1/10$, $\bar{\beta}_2 = 2/5$, i.e., $\alpha_0 = -7/10$, $\alpha_1 = -1/10$ (solid line), $\bar{\beta}_1 = 1/10$, $\bar{\beta}_2 = 1$, i.e., $\alpha_0 = -7/10$, $\alpha_1 = 17/10$ (dashed line), $\bar{\beta}_1 = 1$, $\bar{\beta}_2 = 15$, i.e., $\alpha_0 = 2$, $\alpha_1 = 41$ (dotted line), or $\bar{\beta}_1 = 5$, $\bar{\beta}_2 = 50$, i.e., $\alpha_0 = 14$, $\alpha_1 = 134$ (dot-dashed line); (b) $\text{Im } z$ for $\text{Re } z = 0$ and $\bar{\beta}_1 = \bar{\beta}_2 = 1/10$, i.e., $\alpha_0 = -7/10$, $\alpha_1 = -1$ (solid line), $\bar{\beta}_1 = 1$, $\bar{\beta}_2 = 1/10$, i.e., $\alpha_0 = 2$, $\alpha_1 = -37/10$ (dashed line), $\bar{\beta}_1 = 5$, $\bar{\beta}_2 = 1/10$, i.e., $\alpha_0 = 14$, $\alpha_1 = -157/10$ (dotted line), or $\bar{\beta}_1 = 5$, $\bar{\beta}_2 = 1$, i.e., $\alpha_0 = 14$, $\alpha_1 = -13$ (dot-dashed line).




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8